

# Casgevvy® (exagamglogene autotemcel) (Intravenous)

Document Number: M-0744

Last Review Date: 08/04/2026

Date of Origin: 01/04/2024

Dates Reviewed: 01/2024, 02/2024, 01/2025, 02/2025, 07/2025, 01/2026, 08/2026

## I. Length of Authorization

- Initial: Prior authorization validity will be provided initially for one treatment course (1 dose of Casgevvy).
- Renewal: Prior authorization validity may not be renewed.

## II. Dosing Limits

**Max Units (per dose and over time) [HCPS Unit]:**

- 1 billable unit for one dose

## III. Initial Approval Criteria <sup>1</sup>

Submission of supporting clinical documentation (including but not limited to medical records, chart notes, lab results, and confirmatory diagnostics) related to the medical necessity criteria is REQUIRED on all requests for authorizations. Records will be reviewed at the time of submission as part of the evaluation of this request. Please provide documentation related to diagnosis, step therapy, and clinical markers (i.e., genetic, and mutational testing) supporting initiation when applicable. Please provide documentation via direct upload through the PA web portal or by fax. Failure to submit the medical records may result in the denial of the request due to inability to establish medical necessity in accordance with policy guidelines.

Prior authorization validity is provided in the following conditions:

- Member is at least 2 years of age; **AND**
- Provider has considered use of prophylaxis therapy for seizures with agents other than phenytoin prior to initiating myeloablative conditioning; **AND**
- Member has been screened and found negative for hepatitis B virus (HBV), hepatitis C virus (HCV), and human immunodeficiency virus 1 & 2 (HIV-1/HIV-2) in accordance with clinical guidelines prior to collection of cells (leukapheresis); **AND**
- Provider will confirm that member will not receive live vaccines concurrently while immunosuppressed; **AND**
- Member does not have a history of hypersensitivity to dimethyl sulfoxide (DMSO) or dextran 40; **AND**

- Member has not received other gene therapies used for the treatment of sickle cell disease OR beta thalassemia [e.g., Lyfgenia® (lovotibeglogene autotemcel), Zynteglo® (betibeglogene autotemcel), etc.]; **AND**
- Member will not receive therapy concomitantly with any of the following:
  - Iron chelators for at least 7-days prior to myeloablative conditioning and for 6 months post-treatment for myelosuppressive iron chelators (e.g., deferiprone) OR 3-months post-treatment for non-myelosuppressive iron chelators; **AND**
  - Disease-modifying agents (e.g., hydroxyurea or crizanlizumab) for at least 8-weeks prior to mobilization and conditioning; **AND**
- Member is a candidate for autologous hematopoietic stem cell transplant (HSCT) and has not had prior HSCT; **AND**
- For members under 18 years of age, the member does not have a known and available suitable 10/10 human leukocyte antigen matched related donor willing to participate in an allogeneic HSCT; **AND**

*§ Requests for subsequent use of exagamglogene after receipt of other gene therapies for sickle cell disease or beta thalassemia (e.g., lovotibeglogene, betibeglogene, etc.) will be evaluated on a case-by-case basis*

### **Sickle Cell Disease † Φ<sup>1,3</sup>**

- Member has a confirmed diagnosis of sickle-cell disease with one of the following genotypes βS/βS or βS/β0 or βS/β+ (Note: Additional genotypes will be considered on a case-by-case basis based on disease severity) as determined by one of the following:
  - Identification of significant quantities of hemoglobin S (HbS) with or without an additional abnormal β-globin chain variant by hemoglobin assay; **OR**
  - Identification of biallelic *HBB* pathogenic variants where at least one allele is the p.Glu6Val pathogenic variant on molecular genetic testing; **AND**
- Confirmation that member weighs at least 12 kg prior to the start of mobilization; **AND**
- Member has uncontrolled disease despite treatment with hydroxyurea OR crizanlizumab at any point in the past (*Note: trial of crizanlizumab not applicable to members less than 16 years of age*) OR has experienced intolerance OR has required repeat transfusions to treat symptomatic disease and/or reduce the risk of stroke; **AND**
- Member will be transfused prior to apheresis to a total hemoglobin (Hb) ≤ 11 g/dL and a HbS level < 30% and member will be transfused at least 8 weeks prior to initiation of myeloablative conditioning (with aforementioned Hb and HbS goals); **AND**
- Member will not receive granulocyte-colony stimulating factor (G-CSF) for the mobilization of hematopoietic stem cells (HSC); **AND**

- Member has severe, symptomatic disease despite treatment with supportive care measures, as experienced by one or more of the following:
  - Member has echocardiographic evidence of a tricuspid regurgitant jet velocity (TRJV) of > 2.5 m/s ; **OR**
  - Member has had or has a history of an overt stroke (Note: Defined as a sudden neurologic change lasting more than 24 hours that is accompanied by cerebral MRI changes); **OR**
  - Member has experienced an ‘acute chest syndrome’ episode, defined as an acute event with pneumonia-like symptoms and the presence of a new pulmonary infiltrate in the previous 2 years; **OR**
  - Member experienced two or more vaso-occlusive events/crises (VOE/VOC)\* in the previous year

*\*VOE/VOC is defined as an event requiring a visit to a medical facility for evaluation which results in a diagnosis of such being documented due to one (or more) of the following: acute pain, acute chest syndrome, acute splenic sequestration, acute hepatic sequestration, priapism lasting > 2 hours AND necessitating subsequent interventions such as opioid pain management, non-steroidal anti-inflammatory drugs, RBC transfusion, etc.*

#### **Beta Thalassemia † Φ<sup>1,2,11</sup>**

- Member has a documented diagnosis of homozygous beta thalassemia or compound heterozygous beta thalassemia including β-thalassemia/hemoglobin E (HbE) as outlined by the following:
  - Member diagnosis is confirmed by *HBB* sequence gene analysis showing biallelic pathogenic variants; **OR**
  - Member has severe microcytic hypochromic anemia, absence of iron deficiency, anisopoikilocytosis with nucleated red blood cells on peripheral blood smear, and hemoglobin analysis that reveals decreased amounts or complete absence of hemoglobin A (HbA) and increased HbA<sub>2</sub> with or without increased amounts of hemoglobin F (HbF); **AND**
- Member has transfusion-dependent disease defined as a history of transfusions of at least 100 mL/kg/year or ≥10 units/year of packed red blood cells (pRBCs) in the 2 years preceding therapy; **AND**
- Member will be transfused prior to apheresis to a total Hb ≥ 11 g/dL for 60 days prior to myeloablative conditioning; **AND**
- Member does not have any of the following:

- Severely elevated iron in the heart (i.e., members with cardiac T2\* less than 10 msec by magnetic resonance imaging [MRI] or left ventricular ejection fraction [LVEF] < 45% by echocardiogram); **OR**
- Advanced liver disease [i.e., AST or ALT > 3 times the upper limit of normal (ULN), or direct bilirubin value > 2.5 times the ULN, or if a liver biopsy demonstrated bridging fibrosis or cirrhosis]

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

#### IV. Renewal Criteria <sup>1</sup>

- Duration of authorization has not been exceeded (*refer to Section I*).

#### V. Dosage/Administration <sup>1</sup>

| Indication                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | Dose                                                                                                                                                                                                                                                                                                                                  |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Sickle Cell Disease or Beta Thalassemia                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | <p>Casgevy is provided as a single dose for intravenous infusion containing a suspension of CD34+ cells in one or more vials to achieve the member-specific dose. Administer all vials.</p> <ul style="list-style-type: none"> <li>• The minimum recommended dose of Casgevy is <math>3 \times 10^6</math> CD34+ cells/kg.</li> </ul> |
| <p>- <i>Sickle Cell Disease: Mobilization should occur using single agent plerixafor. Refer to the full Casgevy Prescribing Information for the recommended plerixafor dosing regimen.</i></p> <p>- <i>Beta Thalassemia: Mobilization should occur using both plerixafor and Granulocyte-Colony Stimulating Factor (G-CSF). Refer to the full Casgevy Prescribing Information for the recommended plerixafor and G-CSF dosing regimen.</i></p> <p>- <i>Myeloablative conditioning (e.g., busulfan) should not occur until Casgevy (and back-up cell collection) are received.</i></p> <p>- <i>Consider hepatic veno occlusive disease (VOD)/hepatic sinusoidal obstruction syndrome prophylaxis prior to initiating busulfan conditioning. Refer to the full Casgevy Prescribing Information for the recommended busulfan myeloablative conditioning dosing regimen.</i></p> <p>- <i>Casgevy must be administered between 48 hours and 7 days after the last dose of the myeloablative conditioning regimen.</i></p> <p>- <i>Casgevy is for autologous use only. Before infusion, confirm that the member's identity matches the unique member information located on the Casgevy vial(s). Do not infuse if the information on the member-specific label on any of the vials does not match the intended member.</i></p> |                                                                                                                                                                                                                                                                                                                                       |

#### VI. Billing Code/Availability Information

##### HCPCS Code(s):

- J3392 – Injection, exagamglogene autotemcel, per treatment; 1 billable unit = 1 treatment

##### NDC:

- Casgevy containing a minimum of  $3.0 \times 10^6$  CD34+ cells/kg of body weight, in one or more vials packaged in carton(s): 51167-0290-xx

## VII. References

1. Casgevy [package insert]. Boston, MA; Vertex Pharmaceuticals Incorporated., July 2026. Accessed July 2026.
2. Frangoul H, Altshuler D, Cappellini MD, et al. CRISPR-Cas9 Gene Editing for Sickle Cell Disease and  $\beta$ -Thalassemia. Jan 21, 2021 N Engl J Med 2021; 384:252-260 DOI: 10.1056/NEJMoa2031054. Epub 2020 Dec 5.
3. Bender MA, Carlberg K. Sickle Cell Disease. 2003 Sep 15 [Updated 2025 Feb 13]. In: Adam MP, Bick S, Mirzaa GM, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2025. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1377/>.
4. Yawn BP, Buchanan GR, Afeniyi-Annan AN, et al. Management of sickle cell disease: summary of the 2014 evidence-based report by expert panel members. JAMA. 2014 Sep 10;312(10):1033-48.
5. Tisdale JF, Pierciey FJ, Bonner M, et al. (2020) Safety and feasibility of hematopoietic progenitor stem cell collection by mobilization with plerixafor followed by apheresis vs bone marrow harvest in patients with sickle cell disease in the multi-center HGB-206 trial. Am J Hematol E239–E242. <https://doi.org/10.1002/ajh.25867>.
6. Palmer J, McCune JS, Perales M-A, et al. (2016) Personalizing Busulfan-Based Conditioning: Considerations from the American Society for Blood and Marrow Transplantation Practice Guidelines Committee. Biol Blood Marrow Transplant 1915–1925. <https://doi.org/10.1016/j.bbmt.2016.07.013>
7. Brunson A, Keegan THM, Bang H, et al. (2017) Increased risk of leukemia among sickle cell disease patients in California. Blood 130:1597–1599. doi: 10.1182/blood-2017-05-783233.
8. Seminog OO, Ogunlaja OI, Yeates D, Goldacre MJ (2016) Risk of individual malignant neoplasms in patients with sickle cell disease: English national record linkage study. J R Soc Med 109:303–309. doi: 10.1177/0141076816651037.
9. Brusson M, Miccio A. Genome editing approaches to beta-hemoglobinopathies. Prog Mol Biol Transl Sci. 2021;182:153-183. doi: 10.1016/bs.pmbts.2021.01.025. Epub 2021 Mar 1.
10. Modarai SR, Kanda S, Bloh K, et al. Precise and error-prone CRISPR-directed gene editing activity in human CD34+ cells varies widely among patient samples. Gene Ther. 2021 Feb;28(1-2):105-113. doi: 10.1038/s41434-020-00192-z. Epub 2020 Sep 1.
11. Langer AL. Beta-Thalassemia. 2000 Sep 28 [Updated 2024 Feb 8]. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1426/>.
12. Locatelli F, Thompson AA, Kwiatkowski JL, et al. Betibeglogene Autotemcel Gene Therapy for Non- $\beta$ (0)/ $\beta$ (0) Genotype  $\beta$ -Thalassemia. N Engl J Med. 2022 Feb 3;386(5):415-427. doi: 10.1056/NEJMoa2113206. Epub 2021 Dec 11.

13. Locatelli F, Lang P, Wall D, et al; CLIMB THAL-111 Study Group. Exagamglogene Autotemcel for Transfusion-Dependent  $\beta$ -Thalassemia. N Engl J Med. 2024 May 9;390(18):1663-1676. doi: 10.1056/NEJMoa2309673. Epub 2024 Apr 24. PMID: 38657265.
14. Frangoul H, de la Fuente J, Chopra Y, et al; CLIMB THAL-141 and CLIMB SCD-151 Study Groups. Exa-cel in Children with Transfusion-Dependent  $\beta$ -Thalassemia or Sickle Cell Disease. N Engl J Med. 2026 Jun 11. doi: 10.1056/NEJMoa2603387. Epub ahead of print. PMID: 42274009.
15. Frangoul H, Locatelli F, Sharma A, et al; CLIMB SCD-121 Study Group. Exagamglogene Autotemcel for Severe Sickle Cell Disease. N Engl J Med. 2024 May 9;390(18):1649-1662. doi: 10.1056/NEJMoa2309676. Epub 2024 Apr 24. PMID: 38661449.

## Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime's assessment led to for each.

| Factor                     | Conclusion            |
|----------------------------|-----------------------|
| Indication                 | Yes: Consider for PA  |
| Safety and efficacy        | No: PA not a priority |
| Potential for misuse/abuse | No: PA not a priority |
| Cost of drug               | Yes: Consider for PA  |

## Appendix 1 – Covered Diagnosis Codes

| ICD-10 | ICD-10 Description                               |
|--------|--------------------------------------------------|
| D56.1  | Beta thalassemia                                 |
| D56.5  | Hemoglobin E-beta thalassemia                    |
| D57.00 | Hb-SS disease with crisis, unspecified           |
| D57.01 | Hb-SS disease with acute chest syndrome          |
| D57.02 | Hb-SS disease with splenic sequestration         |
| D57.03 | Hb-SS disease with cerebral vascular involvement |
| D57.04 | Hb-SS disease with dactylitis                    |

|         |                                                                                     |
|---------|-------------------------------------------------------------------------------------|
| D57.09  | Hb-SS disease with crisis with other specified complication                         |
| D57.1   | Sickle-cell disease without crisis                                                  |
| D57.20  | Sickle-cell/Hb-C disease without crisis                                             |
| D57.211 | Sickle-cell/Hb-C disease with acute chest syndrome                                  |
| D57.212 | Sickle-cell/Hb-C disease with splenic sequestration                                 |
| D57.213 | Sickle-cell/Hb-C disease with cerebral vascular involvement                         |
| D57.214 | Sickle-cell/Hb-C disease with dactylitis                                            |
| D57.218 | Sickle-cell/Hb-C disease with crisis with other specified complication              |
| D57.219 | Sickle-cell/Hb-C disease with crisis, unspecified                                   |
| D57.40  | Sickle-cell thalassemia without crisis                                              |
| D57.411 | Sickle-cell thalassemia, unspecified, with acute chest syndrome                     |
| D57.412 | Sickle-cell thalassemia, unspecified, with splenic sequestration                    |
| D57.413 | Sickle-cell thalassemia, unspecified, with cerebral vascular involvement            |
| D57.414 | Sickle-cell thalassemia, unspecified, with dactylitis                               |
| D57.418 | Sickle-cell thalassemia, unspecified, with crisis with other specified complication |
| D57.419 | Sickle-cell thalassemia, unspecified, with crisis                                   |
| D57.42  | Sickle-cell thalassemia beta zero without crisis                                    |
| D57.431 | Sickle-cell thalassemia beta zero with acute chest syndrome                         |
| D57.432 | Sickle-cell thalassemia beta zero with splenic sequestration                        |
| D57.433 | Sickle-cell thalassemia beta zero with cerebral vascular involvement                |
| D57.434 | Sickle-cell thalassemia beta zero with dactylitis                                   |
| D57.438 | Sickle-cell thalassemia beta zero with crisis with other specified complication     |
| D57.439 | Sickle-cell thalassemia beta zero with crisis, unspecified                          |
| D57.44  | Sickle-cell thalassemia beta plus without crisis                                    |
| D57.451 | Sickle-cell thalassemia beta plus with acute chest syndrome                         |
| D57.452 | Sickle-cell thalassemia beta plus with splenic sequestration                        |
| D57.453 | Sickle-cell thalassemia beta plus with cerebral vascular involvement                |
| D57.454 | Sickle-cell thalassemia beta plus with dactylitis                                   |
| D57.458 | Sickle-cell thalassemia beta plus with crisis with other specified complication     |
| D57.459 | Sickle-cell thalassemia beta plus with crisis, unspecified                          |
| D57.80  | Other sickle-cell disorders without crisis                                          |
| D57.811 | Other sickle-cell disorders with acute chest syndrome                               |

#### Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

|         |                                                                           |
|---------|---------------------------------------------------------------------------|
| D57.812 | Other sickle-cell disorders with splenic sequestration                    |
| D57.813 | Other sickle-cell disorders with cerebral vascular involvement            |
| D57.814 | Other sickle-cell disorders with dactylitis                               |
| D57.818 | Other sickle-cell disorders with crisis with other specified complication |
| D57.819 | Other sickle-cell disorders with crisis, unspecified                      |

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A

| Medicare Part B Administrative Contractor (MAC) Jurisdictions |                                                                                             |                                                   |
|---------------------------------------------------------------|---------------------------------------------------------------------------------------------|---------------------------------------------------|
| Jurisdiction                                                  | Applicable State/US Territory                                                               | Contractor                                        |
| E (1)                                                         | CA, HI, NV, AS, GU, CNMI                                                                    | Noridian Healthcare Solutions, LLC                |
| F (2 & 3)                                                     | AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ                                                      | Noridian Healthcare Solutions, LLC                |
| 5                                                             | KS, NE, IA, MO                                                                              | Wisconsin Physicians Service Insurance Corp (WPS) |
| 6                                                             | MN, WI, IL                                                                                  | National Government Services, Inc. (NGS)          |
| H (4 & 7)                                                     | LA, AR, MS, TX, OK, CO, NM                                                                  | Novitas Solutions, Inc.                           |
| 8                                                             | MI, IN                                                                                      | Wisconsin Physicians Service Insurance Corp (WPS) |
| N (9)                                                         | FL, PR, VI                                                                                  | First Coast Service Options, Inc.                 |
| J (10)                                                        | TN, GA, AL                                                                                  | Palmetto GBA                                      |
| M (11)                                                        | NC, SC, WV, VA (excluding below)                                                            | Palmetto GBA                                      |
| L (12)                                                        | DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA) | Novitas Solutions, Inc.                           |
| K (13 & 14)                                                   | NY, CT, MA, RI, VT, ME, NH                                                                  | National Government Services, Inc. (NGS)          |
| 15                                                            | KY, OH                                                                                      | CGS Administrators, LLC                           |