

Opdivo® (nivolumab) (Intravenous)

-E-

Document Number: IC-0541

Date Reviewed: 04/2026

Date of Origin: 07/01/2020

Dates Approved: 07/2020, 10/2020, 12/2020, 02/2021, 03/2021, 06/2021, 07/2021, 10/2021, 01/2022, 04/2022, 07/2022, 10/2022, 01/2023, 04/2023, 07/2023, 10/2023, 12/2023, 01/2024, 04/2024, 08/2024, 11/2024, 11/2024, 01/2025, 02/04/2025, 05/05/2025, 06/05/2025, 09/04/2025, 11/04/2025, 03/03/2026, 06/02/2026

I. Length of Authorization ^{Δ 1,35,43,47,49,50,52-54,65,68,72,73,79,81,82,89,129,130,137,139-141}

- Initial: Prior authorization validity will be provided initially for 6 months (180 days), unless otherwise specified.
 - Use in the treatment of Classical Hodgkin Lymphoma (cHL):
 - ❖ Adult cHL in combination with brentuximab vedotin: Prior authorization validity may be provided up to a maximum of 24 weeks of therapy (8 doses).
 - ❖ Pediatric cHL in combination with brentuximab vedotin-based therapy: Prior authorization validity may be provided up to a maximum of 12 weeks of therapy (4 doses).
 - ❖ Adult and Pediatric cHL in combination with ICE (ifosfamide, carboplatin, etoposide): Prior authorization validity may be provided up to a maximum of 12 weeks of therapy (6 doses).
 - ❖ Adult and Pediatric cHL in combination with AVD (doxorubicin, vinblastine, dacarbazine): Prior authorization validity may be provided up to a maximum of 24 weeks of therapy (12 doses).
 - Use in the treatment of Cutaneous Melanoma:
 - ❖ Cutaneous Melanoma neoadjuvant therapy in combination with ipilimumab: Prior authorization validity may be provided for a maximum of 2 doses.
 - ❖ Cutaneous Melanoma adjuvant treatment in combination with ipilimumab: Prior authorization validity may be provided for a maximum of 4 doses.
 - Merkel Cell Carcinoma neoadjuvant therapy: Prior authorization validity may be provided for up to a maximum of 2 doses.
- Renewal: Prior authorization validity may be renewed every 6 months (180 days) thereafter, unless otherwise specified.
 - Esophageal and Esophagogastric/Gastroesophageal Junction Cancer neoadjuvant or perioperative therapy (adenocarcinoma): Prior authorization validity may be provided for a

maximum of 12 weeks of pre-operative therapy (6 doses), followed by a maximum of 36 weeks (9 doses) of postoperative therapy after surgery.

- Gastric Cancer neoadjuvant or perioperative therapy: Prior authorization validity may be provided for a maximum of 12 weeks of pre-operative therapy (6 doses), followed by a maximum of 36 weeks (9 doses) of postoperative therapy after surgery.
- Non-Small Cell Lung Cancer neoadjuvant treatment followed by optional adjuvant treatment: Prior authorization validity may be provided for a maximum of 4 neoadjuvant doses and 13 adjuvant doses.
- Prior authorization validity may NOT be renewed for the following indications:
 - ❖ Adult cHL in combination with brentuximab vedotin
 - ❖ Pediatric cHL in combination with brentuximab vedotin-based therapy
 - ❖ Adult and Pediatric cHL in combination with ICE (ifosfamide, carboplatin, etoposide)
 - ❖ Adult and Pediatric cHL in combination with AVD (doxorubicin, vinblastine, dacarbazine)
 - ❖ Merkel Cell Carcinoma (neoadjuvant therapy)
 - ❖ Cutaneous Melanoma (neoadjuvant therapy)
 - ❖ Cutaneous Melanoma (adjuvant therapy in combination with ipilimumab)
- Prior authorization validity may be renewed up to a maximum of 1 year of therapy* for the following:
 - ❖ Cutaneous Melanoma (adjuvant therapy single agent)
 - ❖ Esophageal and Esophagogastric/Gastroesophageal Junction Cancer (adjuvant therapy following neoadjuvant chemoradiotherapy)
 - ❖ Squamous Cell Skin Cancer (weight-based 3 mg/kg dosing)
 - ❖ Urothelial Carcinoma (adjuvant therapy)
- Prior authorization validity may be renewed up to a maximum of 2 years of therapy* for the following:
 - ❖ Biliary Tract Cancer (subsequent therapy)
 - ❖ Bone Cancer (in combination with ipilimumab)
 - ❖ Cervical Cancer
 - ❖ Esophageal and Esophagogastric/Gastroesophageal Junction Cancer [first-line therapy, subsequent therapy (excluding single agent use for squamous cell carcinoma)]
 - ❖ Gastric Cancer (first-line therapy)
 - ❖ Kaposi Sarcoma (in combination with ipilimumab)
 - ❖ Renal Cell Carcinoma (in combination with cabozantinib)
 - ❖ Pleural Mesothelioma (first-line/induction therapy in combination with ipilimumab)**

- ❖ Peritoneal Mesothelioma (first-line therapy in combination with ipilimumab)**
- ❖ Non-Small Cell Lung Cancer (in combination with ipilimumab with or without platinum-doublet chemotherapy)
- ❖ Ovarian, Fallopian Tube, and Primary Peritoneal Cancer – Clear Cell Carcinoma of the Ovary
- ❖ Squamous Cell Skin Cancer (flat 240 mg dosing)
- ❖ Vaginal Cancer
- ❖ Vulvar Cancer
- ❖ Urothelial Carcinoma (first line therapy in combination with gemcitabine and cisplatin, followed by single-agent maintenance therapy)

** Including pericardial mesothelioma and tunica vaginalis testis mesothelioma

*Note: The maximum number of doses is dependent on the dosing frequency and duration of therapy. Refer to Section V for exact dosage.		
Dosing Frequency	Maximum length of therapy	Maximum number of doses
2 weeks	1 year	26 doses
	2 years	52 doses
3 weeks	2 years	35 doses
4 weeks	1 year	13 doses
	2 years	26 doses

II. Dosing Limits

Max Units (per dose and over time) [HCPCS Unit]:

Indication	Billable Units (BU)	Per unit time (days)
Biliary, cHL, Cutaneous Melanoma, Gastric, Gestational Trophoblastic Neoplasia (GTN), SCCHN, HCC, Kaposi Sarcoma, Ovarian, Fallopian Tube, & Primary Peritoneal Cancer, RCC, & Soft Tissue Sarcoma	1440 billable units	84 days
Anal, Ampullary Adenocarcinoma, Appendiceal, Bone, CLL/SLL, CNS cancers, CRC, Esophageal and Esophagogastric/Gastroesophageal Junction Cancer, Merkel Cell, PM, PeM, Pericardial Mesothelioma, Tunica Vaginalis Testis Mesothelioma, Squamous Cell Skin, PMBCL, NSCLC, SCLC, Small Bowel Adenocarcinoma, Vulvar Cancer, Vaginal Cancer, & Cervical Cancer	2040 billable units	84 days
Uveal Melanoma	6960 billable units	84 days
Uterine Neoplasms	<i>Initial</i> 340 billable units	14 days x 8 doses
	<i>Maintenance</i>	

	480 billable units	28 days
Urothelial Carcinoma (Bladder Cancer)	<i>Initial</i>	
	360 billable units	21 days x 6 doses
	<i>Maintenance</i>	
	480 billable units	28 days

III. Initial Approval Criteria ¹

Prior authorization validity is provided for the following conditions:

- Member is at least 18 years of age (unless otherwise specified); **AND**

Universal Criteria

- Member has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy, unless otherwise specified ^Δ; **AND**
- Therapy will not be used concomitantly with subcutaneous nivolumab; **AND**

Ampullary Adenocarcinoma ‡ ^{2,178e,281e}

- Member has microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) disease as determined by an FDA-approved or Clinical Laboratory Improvement Amendments (CLIA)-compliant test❖; **AND**
 - Used in combination with ipilimumab if good performance status (ECOG PS 0-1 with good biliary drainage and adequate nutritional intake), intermediate performance status (ECOG PS 2), or poor performance status (ECOG PS 3); **AND**
 - Member has intestinal type disease; **AND**
 - Used as first-line therapy for metastatic disease; **OR**
 - Used as subsequent therapy for disease progression; **OR**
 - Used as a single agent; **AND**
 - Member has intestinal type disease; **AND**
 - Used as first-line therapy for metastatic disease if poor performance status (ECOG PS 3); **OR**
 - Used as subsequent therapy for disease progression if intermediate performance status (ECOG PS 2) or poor performance status (ECOG PS 3)

Anal Carcinoma ‡ ^{2,6,35}

- Member has squamous cell carcinoma; **AND**
 - Used as a single agent as subsequent therapy for metastatic disease

Biliary Tract Cancers (Gallbladder Cancer or Intra-/Extra-Hepatic Cholangiocarcinoma) ‡

2,72,173e

- Member has tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
- Used in combination with ipilimumab; **AND**
- Used as subsequent treatment for progression on or after systemic treatment for unresectable, gross residual (R2), or metastatic disease; **AND**
- Disease is refractory to standard therapies or there are no standard treatment options available; **AND**

- | |
|--|
| <ul style="list-style-type: none">• Use of nivolumab will be restricted to members with a contraindication or intolerance to pembrolizumab |
|--|

Urothelial Carcinoma (Bladder Cancer) † ‡ 1,2,30,51,62,92

- Used as a single agent; **AND**
 - Used for locally advanced or metastatic urothelial carcinoma that progressed during or following platinum-containing chemotherapy* †; **OR**
 - Used as adjuvant therapy; **AND**
 - Member has urothelial carcinoma of the bladder †, male bulbar urethra, prostate with stromal invasion, or upper genitourinary (GU) tract (ureter or renal pelvis) †; **AND**
 - Member is at high risk for disease recurrence**; **OR**
- Used in combination with cisplatin and gemcitabine followed by nivolumab maintenance therapy; **AND**
 - Used as first-line systemic therapy in cisplatin eligible members*; **AND**
 - Member has one of the following diagnoses:
 - Locally advanced, unresectable, or metastatic urothelial carcinoma
 - Muscle invasive bladder cancer with local recurrence or persistent disease in a preserved bladder
 - Metastatic or local bladder cancer recurrence post-cystectomy
 - Primary carcinoma of the urethra; **AND**
 - Clinical stage T3-4, cN1-2 disease or cN1-2 palpable inguinal lymph nodes; **OR**
 - Recurrent or metastatic disease
 - Metastatic upper genitourinary (GU) tract tumors
 - Metastatic urothelial carcinoma of the prostate

* **Note:** 10,51,60,70

- If member was progression free for >12 months after platinum therapy, consider re-treatment with platinum-based therapy if the member is still platinum eligible (see below for cisplatin- or platinum-ineligible comorbidities).
 - Cisplatin-ineligible comorbidities may include the following: CrCl < 60 mL/min, ECOG PS ≥ 2 or KPS ≤ 70%, hearing loss of ≥ 25 decibels (dB) at two contiguous frequencies, grade ≥ 2 peripheral neuropathy, or NYHA Heart Failure class ≥ 3. Carboplatin may be substituted for cisplatin in the metastatic setting for cisplatin-ineligible members such as those with a GFR less than 60 mL/min.
 - Platinum-ineligible comorbidities may include the following: CrCl < 30 mL/min, ECOG PS ≥ 3, grade ≥ 2 peripheral neuropathy, or NYHA Heart Failure class > 3, etc.

** **Note:** 1,62

- High risk for disease recurrence is defined as:
 - ypT2-ypT4a or ypN+ for members who received neoadjuvant platinum-based therapy (excluding urothelial carcinoma of the prostate with stromal invasion); **OR**
 - pT3-pT4a or pN+ for members who did not receive neoadjuvant platinum-based therapy

Bone Cancers ‡ 2,72,173e

- Member has one of the following: Chondrosarcoma, Ewing Sarcoma*, Conventional Chordoma (including chondroid), High-Grade undifferentiated pleomorphic sarcoma (UPS), or Osteosarcoma; **AND**
 - Used in combination with ipilimumab; **AND**
 - Member has tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Member has unresectable or metastatic disease that progressed following prior treatment; **AND**
 - Member has no satisfactory alternative treatment options; **AND**
 - Use of nivolumab will be restricted to members with a contraindication or intolerance to pembrolizumab; **OR**
- Member has Dedifferentiated chondrosarcoma; **AND**
 - Used in combination with sunitinib; **AND**
 - Member has locally advanced or metastatic disease that progressed following prior treatment; **OR**
 - Used in combination with ipilimumab; **AND**
 - Member has tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**

- Member has unresectable or metastatic disease that progressed following prior treatment; **AND**
 - Member has no satisfactory alternative treatment options; **AND**
- Use of nivolumab will be restricted to members with a contraindication or intolerance to pembrolizumab

**Other primary round cell tumors of the bone (eg, CIC::DUX4, BCOR::CCNB3) can be treated like Ewing Sarcoma*

Adult Central Nervous System (CNS) Cancers ‡^{2,5,34,41,42,284e}

- Used in combination with ipilimumab for the treatment of brain metastases in members with BRAF non-specific melanoma

Pediatric Central Nervous System (CNS) Cancers ‡^{2,71,270e}

- Member is ≤ 21 years of age; **AND**
- Member has hypermutant diffuse high-grade glioma; **AND**
- Used for recurrent or progressive disease as a single agent (*excluding oligodendroglioma, IDH-mutant and 1p/19q co-deleted or astrocytoma IDH-mutant*)

Cervical Cancer ‡^{2,49,63,97e,258e}

- Used as subsequent therapy; **AND**
 - Member has recurrent or metastatic adenocarcinoma, adenosquamous carcinoma or squamous cell carcinoma; **AND**
 - Used in combination with ipilimumab; **OR**
 - Used as a single agent; **AND**
 - Tumor expresses PD-L1 (e.g., CPS ≥1) as determined by an FDA-approved or CLIA-compliant test❖; **OR**
 - Member has persistent, recurrent, or metastatic small cell neuroendocrine carcinoma of the cervix (NECC); **AND**
 - Used in combination with ipilimumab; **AND**
- Use of nivolumab with ipilimumab will be restricted to members with a contraindication or intolerance to topotecan or irinotecan

Colorectal Cancer (CRC) ‡^{1,2,31,32,59,106e,107e,178e,285e-287e}

- Member is at least 12 years of age; **AND**
 - Member has microsatellite instability-high (MSI-H)/mismatch repair deficient (dMMR) disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**

- Used in combination with ipilimumab or as a single agent; **AND**
 - Used as primary/initial treatment for unresectable or medically inoperable, recurrent, advanced, or metastatic disease; **OR**
 - Used as subsequent therapy for unresectable or medically inoperable, advanced, or metastatic disease; **AND**
 - For single agent only: Disease progressed following treatment with a fluoropyrimidine-, oxaliplatin-, and/or irinotecan-based chemotherapy, unless contraindicated; **OR**
- Used in combination with ipilimumab; **AND**
 - Used as neoadjuvant therapy for advanced or metastatic disease; **OR**
 - Used for advanced or metastatic disease; **AND**
 - Member has previously received checkpoint inhibitor monotherapy; **OR**
- Member has polymerase epsilon/delta (POLE/POLD1) mutation with ultra-hypermutated phenotype [e.g., tumor mutational burden (TMB) >50 mut/Mb] as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Used as a single agent; **AND**
 - Used as subsequent therapy for locally unresectable or medically inoperable disease, advanced, or metastatic disease

Appendiceal Neoplasms and Cancers ‡^{2,133}

- Member has microsatellite instability-high (MSI-H)/mismatch repair deficient (dMMR) disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Used as a single agent or in combination with ipilimumab if no previous treatment with a checkpoint inhibitor (*Note: Nivolumab + ipilimumab may be considered as subsequent therapy if checkpoint inhibitor monotherapy was previously received*); **AND**
 - Used for recurrent, progressive, metastatic peritoneal-only, or extraperitoneal disease; **OR**
- Member has polymerase epsilon/delta (POLE/POLD1) mutation with ultra-hypermutated phenotype [e.g., tumor mutational burden (TMB) >50 mut/Mb] as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Used as a single agent; **AND**
 - Used for recurrent, progressive, metastatic peritoneal-only, or extraperitoneal disease; **AND**
 - Used as subsequent therapy

Esophageal Cancer and Esophagogastric/Gastroesophageal Junction Cancers † ‡ Φ

1,2,44,52,56,69,133e,155e

- Used as first-line therapy in members with no prior checkpoint inhibitor therapy or no tumor progression while on therapy with a checkpoint inhibitor; **AND**
 - Member has squamous cell carcinoma; **AND**
 - Member is not a surgical candidate or has unresectable advanced, recurrent, or metastatic disease; **AND**
 - Used in combination with ipilimumab OR in combination with fluorouracil or capecitabine AND cisplatin or oxaliplatin; **AND**
 - Tumor expresses PD-L1 (CPS ≥1) as determined by an FDA-approved or CLIA compliant test❖; **AND**

Used in combination with ipilimumab ONLY:

 - Use of nivolumab will be restricted to members with a contraindication or intolerance to one of the following:
 - Nivolumab/(fluorouracil or capecitabine)/(cisplatin or oxaliplatin)
 - Pembrolizumab/(fluorouracil or capecitabine)/(cisplatin or oxaliplatin); **OR**

- Member has adenocarcinoma; **AND**
 - Member is not a surgical candidate or has unresectable advanced, recurrent, or metastatic disease; **AND**
 - Used in combination with oxaliplatin and either fluorouracil or capecitabine for HER2 negative disease; **AND**
 - Tumor expresses PD-L1 (CPS ≥1) as determined by an FDA-approved or CLIA compliant test❖; **OR**
- Used as subsequent therapy in members with no prior checkpoint inhibitor therapy or no tumor progression while on therapy with a checkpoint inhibitor; **AND**
 - Member has squamous cell carcinoma †; **AND**
 - Member is not a surgical candidate or has unresectable advanced, recurrent, or metastatic disease; **AND**
 - Used as a single agent; **AND**
 - Member is refractory or intolerant to at least one prior fluoropyrimidine- and platinum-based regimen; **OR**
- Used as adjuvant/postoperative treatment of completely resected disease; **AND**
 - Member has squamous cell carcinoma or adenocarcinoma; **AND**

- Used as a single agent in members with residual disease following neoadjuvant/preoperative chemoradiotherapy (CRT); **OR**
 - Used as neoadjuvant or perioperative therapy; **AND**
 - Member has microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Member has adenocarcinoma; **AND**
 - Used in combination with ipilimumab; **AND**
 - Used as primary treatment for members who are medically fit for surgery with cT2, N0 (high-risk lesions: lymphovascular invasion, ≥ 3cm, poorly differentiated), cT1b-cT2, N+ or cT3-cT4a, Any N disease; **AND**
- For dMMR disease AND neoadjuvant therapy only:

 - Use of nivolumab with ipilimumab will be restricted to members with a contraindication or intolerance to dostarlimab; **OR**
- Used as a single agent; **AND**
 - Used as postoperative management following R0 resection in members who have received preoperative therapy with nivolumab and ipilimumab

Gastric Cancer † ‡ Φ^{1,2,53,56}

- Used as first-line therapy in members with no prior checkpoint inhibitor therapy or no tumor progression while on therapy with a checkpoint inhibitor; **AND**
 - Member is not a surgical candidate or has unresectable, advanced, recurrent, or metastatic disease; **AND**
 - Used in combination with oxaliplatin and either fluorouracil or capecitabine for HER2 negative disease; **AND**
 - Tumor expresses PD-L1 (CPS ≥1) as determined by an FDA-approved or CLIA compliant test❖; **OR**
 - Member has microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) disease as determined by an FDA-approved or CLIA-compliant test❖ (independent of PD-L1 status); **OR**
- Used as neoadjuvant or perioperative therapy; **AND**
 - Member has microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Used in combination with ipilimumab; **AND**
 - Used as primary treatment prior to surgery for potentially resectable locoregional disease (cT2 or higher, any N) in members who are medically fit for surgery; **AND**

For dMMR disease AND neoadjuvant therapy only:

- Use of nivolumab with ipilimumab will be restricted to members with a contraindication or intolerance to dostarlimab; **OR**
- Used as a single agent; **AND**
 - Used as postoperative management following R0 resection in members who have received preoperative therapy with nivolumab and ipilimumab

Gestational Trophoblastic Neoplasia ‡ ^{2,36, 203e,226e}

- Used in combination with ipilimumab; **AND**
 - Member has multiagent chemotherapy-resistant disease; **AND**
 - Member has intermediate placental site trophoblastic tumor (PSTT) or epithelioid trophoblastic tumor (ETT); **AND**
 - Member has recurrent or progressive disease; **OR**
 - Member has high risk disease (i.e., ≥7 Prognostic score or stage IV disease); **AND**
- Use of nivolumab will be restricted to members with a contraindication or intolerance to avelumab

Squamous Cell Carcinoma of the Head and Neck (SCCHN) † ‡ ^{1,2,29,78}

- Member has Very Advanced Head and Neck Cancer*; **AND**
 - Member has NON-nasopharyngeal cancer; **AND**
 - Used as a single agent; **AND**
 - Member has unresectable, recurrent, persistent, or metastatic disease; **AND**
 - Disease has progressed on or after platinum-containing chemotherapy; **OR**
 - Used in combination with cetuximab; **AND**
 - Member has unresectable, recurrent, persistent, or metastatic disease; **AND**
 - Used as first-line therapy; **AND**

- Use of nivolumab will be restricted to members with a contraindication or intolerance to one of the following regimens:
 - ❖ Pembrolizumab/(cisplatin or carboplatin)/5-FU
 - ❖ Generically available agent/regimen (e.g., cisplatin/paclitaxel, etc. [see *NCCN Head and Neck Cancers guideline for complete list of alternatives*]); **OR**
- Used as subsequent therapy for platinum-resistant or platinum-ineligible disease

** Very Advanced Head and Neck Cancer includes: newly diagnosed (M0) locally advanced T4b, N0-3 disease, newly diagnosed unresectable regional nodal disease, or those unfit for surgery, metastatic disease at initial presentation (M1), or recurrent or persistent disease with or without distant metastases.*

Hepatocellular Carcinoma (HCC) † ‡ Φ 1,2,21,86,87,38e-40e

- Member does not have Child-Turcotte-Pugh (CTP) Class C liver disease; **AND**
 - Used as first-line therapy; **AND**
 - Used in combination with ipilimumab; **AND**
 - Used for one of the following:
 - Unresectable disease
 - Extrahepatic/metastatic disease and is deemed ineligible for resection, transplant, or locoregional therapy; **AND**
- Use of nivolumab will be restricted to members with a contraindication or intolerance to tremelimumab/durvalumab; **OR**
- Used as subsequent therapy; **AND**
 - Used in combination with ipilimumab; **AND**
 - Used for one of the following:
 - Member was previously treated with sorafenib †
 - Member had disease progression on or after systemic therapy and has not previously been treated with anti-CTLA4-based combinations

Adult Classical Hodgkin Lymphoma (cHL) † ‡ Φ 1,2,27,28,54,73,117-118,75e

- Used as a single agent; **AND**
 - Member has relapsed or progressive disease after autologous hematopoietic stem cell transplantation (HSCT) and brentuximab vedotin †; **OR**
 - Used for disease that is refractory to at least 3 prior lines of therapy that includes autologous HSCT †; **OR**
 - Used as palliative subsequent therapy; **AND**
 - Member has relapsed or refractory disease; **AND**
 - Member is not a candidate for high-dose therapy and autologous stem cell rescue (HDT/ASCR); **OR**
 - Used post-allogeneic hematopoietic cell transplant; **OR**
- Used in combination with ICE (ifosfamide, carboplatin, etoposide); **AND**
 - Used as subsequent therapy for relapsed or primary refractory disease; **OR**

- Used in combination with brentuximab vedotin; **AND**
 - Used as subsequent therapy for relapsed or primary refractory disease; **OR**
 - Used as primary treatment for members who are not candidates for anthracycline therapy; **AND**
 - Used in combination with or without ISRT; **OR**
- Used in combination with doxorubicin, vinblastine, and dacarbazine (AVD); **AND**
 - Used as primary treatment for stage III-IV disease

Pediatric Classical Hodgkin Lymphoma (cHL) † ‡ 1,2,27,28,55,117-118,131-132

- Member is ≤ 18 years of age* unless otherwise specified; **AND**
 - Used as primary treatment for stage III-IV disease; **AND**
 - Used in combination with doxorubicin, vinblastine and dacarbazine (AVD) (*applies to members ≥12 years of age ONLY*); **OR**
 - Member has relapsed or refractory disease; **AND**
 - Used in combination with ifosfamide, carboplatin, etoposide (ICE)**; **OR**
 - Used as a single agent**; **AND**
 - Member is heavily pretreated with platinum or anthracycline-based chemotherapy or a decrease in cardiac function was observed; **OR**
 - Used in combination with brentuximab vedotin with or without bendamustine; **AND**
 - Member is heavily pretreated with platinum or anthracycline-based chemotherapy or a decrease in cardiac function was observed; **AND**
 - Used as re-induction therapy or as subsequent therapy (if not previously used); **OR**
 - Used as re-induction therapy in highly favorable members who may avoid autologous stem cell rescue (ASCR) (i.e., initial stage other than IIIB or IVB, no prior exposure to RT, duration of CR1 >1 year, absence of extranodal disease or B symptoms at relapse); **AND**
 - ◆ Used in combination with involved-site radiation therapy (ISRT)

* Pediatric Classic Hodgkin Lymphoma may be applicable to adolescent and young adult (AYA) members up to the age of 39 years

**There is no pediatric data for this regimen

Kaposi Sarcoma † 2,79,227e

- Used in combination with ipilimumab; **AND**
- Used as subsequent therapy; **AND**

- Used for relapsed/refractory advanced T1, extensive T0 cutaneous, or nodal disease; **AND**
- Disease has progressed on or not responded to first-line therapy; **AND**
- Disease has progressed on alternate first-line therapy

Renal Cell Carcinoma (RCC) † ‡ 1,2,25,26,66e,164e,202e

- Used in combination with ipilimumab; **AND**
 - Member has clear cell histology; **AND**
 - Used as first-line therapy; **AND**
 - Member has poor or intermediate risk advanced disease †; **OR**
 - Member has relapsed or stage IV disease*; **OR**
 - Used as subsequent therapy (without history of prior immuno-oncology therapy) in members with relapsed or stage IV disease; **OR**
- Used as a single agent; **AND**
 - Used as subsequent therapy in members with advanced, relapsed, or stage IV disease and clear cell histology; **OR**
- Used in combination with cabozantinib (Cabometyx only); **AND**
 - Member has clear cell histology; **AND**
 - Used as first-line therapy for advanced, relapsed, or stage IV disease*; **OR**
 - Member has non-clear cell histology; **AND**
 - Member has relapsed or stage IV disease*

**When used as first-line therapy for stage IV disease, disease must be M1 or unresectable T4, M0*

Cutaneous Melanoma † ‡ Φ 1,2,15-18,82,93,14e,150e-152e

- Used as first-line therapy for unresectable or metastatic* disease; **AND**
 - Member is at least 12 years of age; **AND**
 - Used as a single agent or in combination with ipilimumab; **OR**
- Used as subsequent therapy for unresectable or metastatic* disease; **AND**
 - Member is at least 12 years of age; **AND**
 - Used as re-induction therapy in members who experienced disease control (i.e., complete or partial response or stable disease) and no residual toxicity from prior nivolumab or nivolumab/ipilimumab, but subsequently have disease progression/relapse > 3 months after treatment discontinuation; **AND**
 - Used as a single agent or in combination with ipilimumab; **OR**

- Used after disease progression, intolerance, and/or projected risk of progression with BRAF-targeted therapy; **AND**
 - Used as a single agent or in combination with ipilimumab; **OR**
- Used as adjuvant treatment; **AND**
 - Used as a single agent; **AND**
 - Member is at least 12 years of age; **AND**
 - Member has stage IIB, stage IIC, or metastatic disease and has undergone complete resection †; **OR**
 - Member has stage III disease; **AND**
 - Member has undergone complete resection †; **OR**
 - Member has resected sentinel node positive disease, during radiographic surveillance OR after complete lymph node dissection (CLND); **OR**
 - Member has clinically positive node(s) following wide excision of the primary tumor and therapeutic lymph node dissection (TLND); **OR**
 - Member has clinical satellite/in-transit metastases and has no evidence of disease (NED) after complete excision to clear margins OR NED after initial treatment with local or regional therapy; **OR**
 - Used following wide excision alone or wide excision with negative sentinel lymph node biopsy (*stage IIIB/C/D disease only*); **OR**
 - Used for disease that is sentinel lymph node negative or sentinel lymph node biopsy not performed (*stage IIIB/C/D disease only*); **OR**
 - Member has local satellite/in-transit recurrence and has NED after complete excision OR NED after initial treatment with local or regional therapy; **OR**
 - Member has resectable disease limited to nodal recurrence following excision of the recurrence; **OR**
 - Member has oligometastatic disease and NED following metastasis-directed therapy (i.e., T-VEC/intralesional therapy, stereotactic ablative radiation therapy or complete resection) OR following systemic therapy followed by resection; **OR**
 - Used in combination with ipilimumab; **AND**
 - Member has oligometastatic disease and no evidence of disease following metastasis-directed therapy (i.e., complete resection, stereotactic ablative radiation therapy or T-VEC/intralesional therapy) OR following systemic therapy followed by resection; **OR**
 - Used as neoadjuvant therapy; **AND**
 - Used in combination with ipilimumab; **AND**
 - Member has stage III disease; **AND**

- Used as primary treatment for clinically positive, resectable nodal disease; **OR**
- Used for limited resectable disease with clinical satellite/in-transit metastases; **OR**
- Member has limited resectable local satellite/in-transit recurrence; **OR**
- Member has resectable disease limited to nodal recurrence

**Metastatic disease includes stage III unresectable/borderline resectable disease with clinically positive node(s) or clinical satellite/in-transit metastases, as well as unresectable/borderline resectable local satellite/in-transit recurrence, unresectable nodal recurrence, oligometastatic disease, and widely disseminated distant metastatic disease and/or brain metastases.*

Uveal Melanoma ‡ 2,19,20,80

- Member has metastatic or unresectable disease; **AND**
- Used in combination with ipilimumab

Merkel Cell Carcinoma ‡ 2,4,33,65,189e

- Used as neoadjuvant treatment; **AND**
 - Used as a single agent; **AND**
 - Member is a surgical candidate with primary clinical N0 locally advanced disease where curative surgery and curative radiation therapy were originally deemed not feasible; **OR**
 - Member has primary clinical N+, M0 regional disease with biopsy positive draining nodal basin; **OR**
- Used for M1 disseminated disease; **AND**
 - Used as a single agent; **AND**

First-line therapy ONLY:

- Use of nivolumab will be restricted to members with a contraindication or intolerance to retifanlimab; **OR**

- Used in combination with ipilimumab; **AND**
 - Used for disease that is refractory to PD-1 therapy; **OR**
- Used for recurrent in-transit N+ regional disease; **AND**
 - Curative surgery and curative radiation therapy (RT) are not feasible; **AND**
 - Used as a single agent; **AND**

First-line therapy ONLY:

- Use of nivolumab will be restricted to members with a contraindication or intolerance to retifanlimab; **OR**

- Used in combination with ipilimumab; **AND**
 - Used for disease that is refractory to PD-1 therapy; **OR**
- Used for recurrent N+ regional disease; **AND**
 - Curative surgery and curative radiation therapy (RT) are not feasible; **AND**
 - Used as a single agent; **AND**

First-line therapy ONLY:

- Use of nivolumab will be restricted to members with a contraindication or intolerance to retifanlimab; **OR**

- Used in combination with ipilimumab; **AND**
 - Used for disease that is refractory to PD-1 therapy

Peritoneal Mesothelioma (PeM)* ‡^{2,64,90}

- Used as a single agent or in combination with ipilimumab as subsequent therapy (if platinum chemotherapy was administered first-line); **OR**
- Used in combination with ipilimumab as first-line therapy; **AND**
 - Used as adjuvant treatment for medically operable disease following cytoreductive surgery (CRS) and hyperthermic intraperitoneal chemotherapy (HIPEC); **AND**
 - Member has surgical or pathologic high-risk features**; **OR**
 - Member has medically inoperable disease and/or complete cytoreduction not achievable, or presence of any high-risk features**; **OR**
 - Member has disease progression following CRS + HIPEC if no prior adjuvant systemic therapy was given

**Note: May also be used for pericardial mesothelioma and tunica vaginalis testis mesothelioma.*

***High-risk features include: biphasic/sarcomatoid histology, nodal metastasis, Ki-67 >9%, thrombocytosis, PS=2, bicavitary disease, high disease burden/incomplete cytoreduction (Peritoneal Cancer Index [PCI] >17, completeness of cytoreduction (cc) score >1)*

Pleural Mesothelioma (PM)* † ‡ Φ^{1,2,37,38,47,64,81}

- Used as a single agent or in combination with ipilimumab as subsequent therapy (if platinum chemotherapy was administered first-line); **OR**
- Used in combination with ipilimumab; **AND**
 - Used as first-line therapy; **AND**
 - Disease is medically inoperable or unresectable; **OR**
 - Used as induction therapy prior to surgical exploration; **AND**

- Member has clinical stage I disease and epithelioid histology

**Note: May also be used for pericardial mesothelioma and tunica vaginalis testis mesothelioma.*

Non-Small Cell Lung Cancer (NSCLC) † ‡ 1,2,11,22,23,43,45,46,120,43e-45e,51e-53e,56e,123e,124e,163e,186e-188e

- Member has resectable (tumors $\geq$ 4 cm or node positive) disease; **AND**
 - Member has no known EGFR mutations or ALK gene fusions; **AND**
 - Used as neoadjuvant therapy in combination with platinum-doublet chemotherapy (e.g., cisplatin/carboplatin in combination with paclitaxel, pemetrexed, gemcitabine, or docetaxel) with the option of continuing single-agent nivolumab as adjuvant treatment after surgery; **OR**
- Used for recurrent, advanced, or metastatic disease; **AND**
 - Used as first-line therapy; **AND**
 - Used for one of the following:
 - Members who have tumors that are negative for actionable biomarkers** (may be KRAS G12C mutation positive)
 - Members who are positive for one of the following biomarkers: EGFR exon 20, KRAS G12C, BRAF V600E, NTRK1/2/3 gene fusions, MET exon 14 skipping, ERBB2 (HER2), or NRG1 gene fusion; **AND**
 - Used in combination with one of the following:
 - Ipilimumab
 - Ipilimumab and platinum-doublet chemotherapy (e.g., pemetrexed and either carboplatin or cisplatin for nonsquamous cell histology, or paclitaxel and carboplatin for squamous cell histology, etc.); **AND**

Squamous NSCLC:

- Use of nivolumab in combination with ipilimumab (with or without platinum-doublet chemotherapy) will be restricted to members with a contraindication or intolerance to cemiplimab/paclitaxel/(carboplatin or cisplatin); **OR**

Nonsquamous NSCLC:

- Use of nivolumab in combination with ipilimumab (with or without platinum-doublet chemotherapy) will be restricted to members with a contraindication or intolerance to cemiplimab/(paclitaxel or pemetrexed)/(carboplatin or cisplatin); **OR**

- Used as subsequent therapy; **AND**
 - Used as a single agent; **OR**
 - Used for one of the following:

- Members who are positive for one of the following biomarkers and have received prior targeted therapy§: EGFR S768I, L861Q, and/or G719X
- Members who are positive for one of the following biomarkers: BRAF V600E, NTRK1/2/3 gene fusions, MET exon 14 skipping, or ERBB2 (HER2); **AND**
- Used in combination with one of the following:
 - Ipilimumab
 - Ipilimumab and platinum-doublet chemotherapy (e.g., pemetrexed and either carboplatin or cisplatin for nonsquamous cell histology, or paclitaxel and carboplatin for squamous cell histology, etc.); **AND**

Squamous NSCLC:

- Use of nivolumab in combination with ipilimumab (with or without platinum-doublet chemotherapy) will be restricted to members with a contraindication or intolerance to cemiplimab/paclitaxel/(carboplatin or cisplatin); **OR**

Nonsquamous NSCLC:

- Use of nivolumab in combination with ipilimumab (with or without platinum-doublet chemotherapy) will be restricted to members with a contraindication or intolerance to cemiplimab/(paclitaxel or pemetrexed)/(carboplatin or cisplatin); **OR**

- Used as continuation maintenance therapy in combination with ipilimumab; **AND**
 - Member has achieved a response or stable disease following first-line therapy with nivolumab and ipilimumab with or without chemotherapy

*** Note: Actionable biomarkers include EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, NRG1, and ERBB2 (HER2). Complete biomarker testing including molecular assessment of EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, NRG1, and ERBB2 (HER2) via biopsy and/or plasma testing. If a clinically actionable marker is found, it is reasonable to start therapy based on the identified marker. Treatment is guided by available results and, if unknown, these members are treated as though they do not have driver oncogenes.*

§ Genomic Aberration/Mutational Driver Targeted Therapies: Refer to guidelines for appropriate use.

Pediatric Aggressive Mature B-Cell Lymphomas – Primary Mediastinal Large B-Cell Lymphoma (PMBCL) ‡^{2,74-76}

- Member is ≤ 18 years of age*; **AND**
- Used in combination with brentuximab vedotin; **AND**
 - Used for relapsed or refractory disease; **AND**

- Used after autologous stem-cell transplant OR if ineligible for autologous stem-cell transplant, used after 2 or more prior lines of therapy; **OR**
- Used as consolidation/additional therapy if a partial response was achieved after therapy for relapsed or refractory disease

** Pediatric Primary Mediastinal Large B-Cell Lymphoma may be applicable to adolescent and young adult (AYA) members <39 years who are treated in a pediatric oncology setting.*

Small Bowel Adenocarcinoma ‡^{2,31,39,59,191e}

- Used as a single agent or in combination with ipilimumab; **AND**
- Member has microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Member has advanced or metastatic disease (*Note: Nivolumab + ipilimumab may be considered as subsequent therapy if checkpoint inhibitor monotherapy was previously received*); **OR**
 - Member has locally unresectable or medically inoperable disease; **AND**
 - Used as primary treatment; **OR**
- Member has polymerase epsilon/delta (POLE/POLD1) mutation with ultra-hypermutated phenotype [e.g., tumor mutational burden (TMB) > 50 mut/Mb] as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Used as single agent; **AND**
 - Used as subsequent therapy for advanced or metastatic disease

Small Cell Lung Cancer (SCLC) ‡ Φ^{2,24,61,147e}

- Used as subsequent systemic therapy as a single agent for relapsed or progressive disease; **AND**
- Member must demonstrate an inadequate response to topotecan or irinotecan, unless there is a contraindication or intolerance, prior to approval of nivolumab

Soft Tissue Sarcoma ‡^{2,72,84,233e,239e}

- Extremity/Body Wall* or Head/Neck*
 - Used as subsequent therapy for advanced/metastatic disease with disseminated metastases; **AND**
 - Member has myxofibrosarcoma, undifferentiated pleomorphic sarcoma (UPS), dedifferentiated liposarcoma, or undifferentiated sarcomas; **AND**
 - Used in combination with ipilimumab; **AND**

– Use of nivolumab in combination with ipilimumab will be restricted to members with a contraindication or intolerance to pembrolizumab; **OR**

- Member has cutaneous angiosarcoma; **AND**
 - Used in combination with ipilimumab
- Retroperitoneal/Intra-Abdominal**
 - Used as one of the following:
 - Alternative systemic therapy for unresectable or progressive disease after initial therapy for unresectable localized disease; **OR**
 - Palliative subsequent therapy for stage IV disease with disseminated metastases; **AND**
 - Used for one of the following:
 - Member has myxofibrosarcoma, undifferentiated pleomorphic sarcoma (UPS), dedifferentiated liposarcoma, or undifferentiated sarcomas; **AND**
 - Used in combination with ipilimumab; **AND**

– Use of nivolumab in combination with ipilimumab will be restricted to members with a contraindication or intolerance to pembrolizumab; **OR**

- Member has cutaneous angiosarcoma; **AND**
 - Used in combination with ipilimumab

- Angiosarcoma

- Used in combination with ipilimumab; **AND**

○ Member must demonstrate an inadequate response to a generically available agent/regimen (e.g., paclitaxel, doxorubicin, etc. [see *NCCN Soft Tissue Sarcoma guideline for complete list of alternatives*]), unless there is a contraindication or intolerance prior to approval of nivolumab; **OR**

- Dedifferentiated Liposarcoma with or without Concurrent Well-Differentiated Liposarcoma

- Used as in combination with ipilimumab; **AND**
 - Used as subsequent therapy; **AND**

○ Use of nivolumab in combination with ipilimumab will be restricted to members with a contraindication or intolerance to pembrolizumab

**For atypical lipomatous tumor/well-differentiated liposarcoma (ALT/WDLPS) of the extremity, abdominal wall, trunk that was initially diagnosed as ALT/WDLPS and shows evidence of de-differentiation, treat as other soft tissue sarcomas.*

***For well-differentiated liposarcoma (WDLPS-retroperitoneum, paratesticular) with or without evidence of de-differentiation, treat as other soft tissue sarcomas*

Uterine Neoplasms ‡^{2,48,72,173e}

- Member has Endometrial Carcinoma; **AND**
 - Used as subsequent therapy as one of the following:
 - Single agent for recurrent disease that is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) disease as determined by an FDA-approved or CLIA-compliant test❖; **OR**
 - In combination with ipilimumab for recurrent unresectable or metastatic disease that is tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Disease has progressed following prior treatment and there are no satisfactory alternative treatment options; **AND**
 - Will not be used for either of the following:
 - Therapy for locoregional recurrence in members with no prior radiation therapy to site of recurrence, or previous vaginal brachytherapy only; **OR**
 - Therapy after surgical exploration for locoregional recurrence in members with disease confined to the vagina or paravaginal soft tissue; **AND**

Used in combination with ipilimumab ONLY:

- Use of nivolumab will be restricted to members with a contraindication or intolerance to pembrolizumab; **OR**
- Member has Uterine Sarcoma; **AND**
 - Used as subsequent therapy in combination with ipilimumab; **AND**
 - Member has unresectable or metastatic disease that is tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] disease as determined by an FDA-approved or CLIA-compliant test❖; **AND**
 - Disease has progressed following prior treatment and there are no satisfactory alternative treatment options; **AND**

- Use of nivolumab will be restricted to members with a contraindication or intolerance to pembrolizumab

Vulvar Cancer ‡^{2,49,136}

- Used as subsequent treatment for advanced, recurrent, or metastatic disease; **AND**
 - Used as single agent for HPV-related tumors; **OR**

- Used in combination with ipilimumab

Vaginal Cancer ‡^{2,49,97}

- Used as subsequent therapy for recurrent or metastatic disease; **AND**
 - Used in combination with ipilimumab; **OR**
 - Used as a single agent; **AND**
 - Tumor expresses PD-L1 (e.g., CPS ≥1) as determined by an FDA-approved or CLIA-compliant test❖

Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL) ‡^{2,216e}

- Member has histologic transformation (Richter); **AND**
- Used in combination with ibrutinib; **AND**
 - Used as additional therapy for partial response, refractory disease, or progression while on prior treatment ^; **OR**
 - Used as first-line treatment for Richter transformation if previously treated for CLL; **OR**
 - Used as continuation therapy for complete response until progression

^Prior treatment could have included immune checkpoint inhibitor therapy (e.g., PD-1/PD-L1-directed therapy)

Squamous Cell Skin Cancer ‡^{2,129,130}

- Used as a single agent; **AND**

- | |
|---|
| <ul style="list-style-type: none"> • Use of nivolumab will be restricted to members with a contraindication or intolerance to cemiplimab; AND |
|---|

- Member has locally advanced disease; **AND**
 - Used as primary treatment if curative surgery and curative radiation therapy (RT) are not feasible; **OR**
- Member has regional disease that is unresectable, inoperable, or incompletely resected if curative RT is not feasible; **AND**
 - Used as first-line therapy; **OR**
- Member has satellitosis/in-transit metastasis that is unresectable or incompletely resected; **AND**
 - Used as first-line therapy; **OR**
- Member has regional recurrence or distant metastatic disease; **AND**
 - Used as first-line therapy

Ovarian, Fallopian Tube, and Primary Peritoneal Cancer ‡^{2,141}

- Used in combination with ipilimumab; **AND**
 - Member has clear cell carcinoma of the ovary; **AND**
 - Used for persistent or recurrent platinum-resistant disease; **OR**
 - Member has small cell carcinoma of the ovary (hypercalcemic type); **AND**
 - Used for progressive or recurrent disease

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

Enhanced Oncology Value (EOV) Program – Redacted indications

Uses not listed above have inadequate data to support efficacy and are excluded from prior authorization validity.

Other treatment options including, but are not limited to, the following may be appropriate: radiation therapy, surgery, traditional chemotherapy (e.g., platinum, taxane), compassionate use/expanded access programs, clinical trials, supportive care, integrative and complementary therapies.

❖ *If confirmed using an FDA approved assay – <http://www.fda.gov/CompanionDiagnostics>*

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

IV. **Renewal Criteria** ^{△ 1,2,4-6,15-42,43,47,49,50,52-54,68,72,73,79,81,82,89}

Prior authorization validity may be renewed based upon the following criteria:

- Member continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Duration of authorization has not been exceeded (*refer to Section I*); **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe infusion-related reactions, complications of allogeneic hematopoietic stem cell

transplantation (HSCT), severe immune-mediated adverse reactions (e.g., pneumonitis, colitis, hepatitis/hepatotoxicity, endocrinopathies, nephritis/renal dysfunction, adverse skin reactions/rash, etc.), etc.; **AND**

- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread

^Δ Notes:

- Members responding to therapy who relapse ≥ 6 months after discontinuation due to duration (i.e., receipt of 24 months of therapy) are eligible to re-initiate PD-directed therapy.
- Members previously presenting with aggressive disease who are exhibiting stable disease on treatment as their best response (or if therapy improved performance status) may be eligible for continued therapy beyond the 24-month limit without interruption or discontinuation.
- Members who complete adjuvant therapy and progress ≥ 6 months after discontinuation are eligible to re-initiate PD-directed therapy for metastatic disease.
- Members whose tumors, upon re-biopsy, demonstrate a change in actionable mutation (e.g., MSS initial biopsy; MSI-H subsequent biopsy) may be eligible to re-initiate PD-directed therapy and will be evaluated on a case-by-case basis.

V. Dosage/Administration ^Δ 1,4-6,19,20,27,24,31-42,48-50,52-54,55,58,59,61,65,67,68,71-87,89,91,93,96,98-119,121-124,127-137,139-141

Indication	Dose
Ampullary Adenocarcinoma	<u>Single agent:</u> • Administer 3 mg/kg intravenously every 2 weeks, 240 mg intravenously every 2 weeks, or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>In combination with ipilimumab:</u> • Administer 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then 3 mg/kg every 2 weeks, or 240 mg every 2 weeks, or 480 mg every 4 weeks until disease progression or unacceptable toxicity
Anal Carcinoma	<u>Single agent:</u> • Administer 240 mg intravenously every 2 weeks, 480 mg intravenously every 4 weeks, 3 mg/kg intravenously every 2 weeks, or 6 mg/kg intravenously every 4 weeks until disease progression or unacceptable toxicity
Biliary Tract Cancers	<u>Subsequent therapy:</u> • Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity for up to 2 years

Urothelial Carcinoma (Bladder Cancer)	<u>First-line therapy:</u> - Administer 360 mg intravenously every 3 weeks for up to 6 cycles (given in combination with gemcitabine and cisplatin), followed by a single-agent maintenance dose of 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity for up to 2 years <u>Disease progression or second-line treatment:</u> - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>Adjuvant treatment:</u> - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease recurrence or unacceptable toxicity for up to 1 year
Bone Cancer	<u>In combination with ipilimumab:</u> - Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity for up to 2 years <u>Single agent or in combination with sunitinib:</u> - Administer 3 mg/kg intravenously every 2 weeks until disease progression or unacceptable toxicity
Adult CNS Cancers	<u>In combination with ipilimumab:</u> Administer 1 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then 3 mg/kg intravenously every 2 weeks, or 240 mg intravenously every 2 weeks, or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity
Pediatric CNS Cancers	Administer 3 mg/kg intravenously every 2 weeks until disease progression or unacceptable toxicity
Colorectal Cancer (CRC)	<u>Adult members and for pediatric members ≥ 12 years and ≥ 40 kg:</u> Single agent: Administer 3 mg/kg intravenously every 2 weeks, or 240 mg intravenously every 2 weeks, or 480 mg intravenously every 4 weeks, or 6 mg/kg intravenously every 4 weeks until disease progression or unacceptable toxicity In combination with ipilimumab: - Administer 240 mg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), followed by single agent regimen until disease progression or unacceptable toxicity; OR - Administer 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), followed by single agent regimen until disease progression or unacceptable toxicity <u>Pediatric members ≥ 12 years and < 40 kg:</u> Single agent: Administer 3 mg/kg intravenously every 2 weeks or 6 mg/kg every 4 weeks until disease progression or unacceptable toxicity

	• In combination with ipilimumab: Administer 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), followed by single agent regimen until disease progression or unacceptable toxicity.
Appendiceal Neoplasms and Cancers	• Single agent: Administer 3 mg/kg intravenously every 2 weeks, or 240 mg intravenously every 2 weeks, or 480 mg intravenously every 4 weeks, or 6 mg/kg intravenously every 4 weeks until disease progression or unacceptable toxicity • In combination with ipilimumab: Administer 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then follow with the single agent regimen until disease progression or unacceptable toxicity
Esophageal and Esophagogastric/Gastroesophageal Junction Cancer	<u>First line therapy (PD-L1 squamous cell carcinoma)</u> • Administer 240 mg intravenously every 2 weeks or 360 mg intravenously every 3 weeks or 480 mg intravenously every 4 weeks (given in combination with fluoropyrimidine- and platinum-containing chemotherapy) until disease progression or unacceptable toxicity for up to 2 years; OR • Administer 3 mg/kg intravenously every 2 weeks or 360 mg intravenously every 3 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity for up to 2 years <u>First line therapy (PD-L1 adenocarcinoma)</u> • Administer 240 mg intravenously every 2 weeks or 360 mg intravenously every 3 weeks (given in combination with fluoropyrimidine- and platinum-containing chemotherapy) until disease progression or unacceptable toxicity for up to 2 years <u>Subsequent therapy (squamous cell carcinoma):</u> • Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks as a single agent until disease progression or unacceptable toxicity <u>Neoadjuvant/perioperative therapy (MSI-H/dMMR adenocarcinoma):</u> • Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) for 12 weeks, followed by surgery and then post-operative therapy (<i>See below</i>) <u>Post-operative therapy (MSI-H/dMMR adenocarcinoma):</u> • Administer 480 mg intravenously every 4 weeks for 36 weeks (9 doses) <u>Adjuvant therapy following neoadjuvant chemoradiotherapy (adenocarcinoma and squamous carcinoma):</u> • Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity for up to 1 year
Gastric Cancer	<u>First-line therapy:</u>

	- Administer 240 mg intravenously every 2 weeks or 360 mg intravenously every 3 weeks (give in combination with oxaliplatin and either fluorouracil or capecitabine) until disease progression or unacceptable toxicity for up to 2 years <u>Neoadjuvant/perioperative therapy:</u> - Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) for 12 weeks, followed by surgery and then post-operative therapy (<i>See below</i>) <u>Post-operative therapy:</u> - Administer 480 mg intravenously every 4 weeks for 36 weeks (9 doses)
Gestational Trophoblastic Neoplasia (GTN)	<u>In combination with ipilimumab:</u> Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity
SCCHN	<u>Single agent:</u> - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>In combination with cetuximab:</u> - Administer 240 mg intravenously every 2 weeks until disease progression or unacceptable toxicity
Hepatocellular Carcinoma (HCC)	<u>In combination with ipilimumab:</u> Administer 1 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity for up to 2 years
Adult cHL	<u>Single agent:</u> - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>In combination with brentuximab vedotin</u> - Administer 3 mg/kg intravenously every 3 weeks for up to 24 weeks (8 doses) <u>In combination with ICE (ifosfamide, carboplatin, and etoposide)</u> - Administer 240 mg intravenously every 2 weeks for up to 12 weeks (6 doses) <u>In combination with AVD (doxorubicin, vinblastine, dacarbazine)</u> - Administer 240 mg intravenously every 2 weeks for up to 24 weeks (12 doses)
Pediatric cHL	<u>Single agent:</u> - Administer 3 mg/kg intravenously every 2 weeks until disease progression or unacceptable toxicity (<i>Note: There is no pediatric data for this regimen</i>) <u>In combination with brentuximab vedotin-based therapy</u> - Administer 3 mg/kg intravenously every 3 weeks for up to 12 weeks (4 doses) <u>In combination with ICE (ifosfamide, carboplatin, and etoposide)</u> - Administer 240 mg intravenously every 2 weeks for up to 12 weeks (6 doses) (<i>Note: There is no pediatric data for this regimen</i>)

	<u>In combination with AVD (doxorubicin, vinblastine, dacarbazine)</u> - Members ≥ 12 years of age weighing 40 kg or more: Administer 240 mg intravenously every 2 weeks for up to 24 weeks (12 doses) - Members 12 to 17 years of age weighing less than 40 kg: Administer 3 mg/kg (up to 240 mg max) every 2 weeks for up to 24 weeks (12 doses)
Kaposi Sarcoma	<u>In combination with ipilimumab:</u> Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity for up to 2 years
Renal Cell Carcinoma (RCC)	<u>Single agent:</u> - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>In combination with ipilimumab:</u> - Administer 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then follow with the single agent regimen until disease progression or unacceptable toxicity <u>In combination with cabozantinib (Cabometyx):</u> - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity for up to 2 years
Pleural Mesothelioma (PM) & Peritoneal Mesothelioma (PeM) (including pericardial mesothelioma and tunica vaginalis testis mesothelioma)	<u>Single agent:</u> - Administer 3 mg/kg intravenously or 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>In combination with ipilimumab:</u> - Subsequent therapy - Administer 3 mg/kg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity; OR - Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity - All other lines of therapy: - Administer 360 mg intravenously every 3 weeks or 3 mg/kg every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity for up to 2 years
Cutaneous Melanoma	<u>Adult members and pediatric members ≥ 12 years and ≥ 40 kg:</u> <u>Single agent</u> <u>Unresectable or metastatic disease:</u> Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity

	• <u>Adjuvant treatment:</u> Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease recurrence or unacceptable toxicity for up to 1 year In combination with ipilimumab • <u>Unresectable or metastatic disease:</u> Administer 1 mg/kg intravenously or 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then follow with the single agent regimen • <u>Adjuvant treatment:</u> Administer 1 mg/kg intravenously or 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day) • <u>Neoadjuvant treatment:</u> Administer 3 mg/kg intravenously every 3 weeks for up to 2 doses (given in combination with ipilimumab on the same day) <u>Pediatric members ≥ 12 years and < 40 kg:</u> Single agent • <u>Unresectable or metastatic disease:</u> Administer 3 mg/kg intravenously every 2 weeks or 6 mg/kg intravenously every 4 weeks until disease progression or unacceptable toxicity • <u>Adjuvant treatment:</u> Administer 3 mg/kg intravenously every 2 weeks or 6 mg/kg intravenously every 4 weeks until disease recurrence or unacceptable toxicity for up to 1 year In combination with ipilimumab • <u>Unresectable or metastatic disease:</u> Administer 1 mg/kg intravenously or 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then follow with the single agent regimen • <u>Adjuvant treatment:</u> Administer 1 mg/kg intravenously or 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day)
Uveal Melanoma	<u>In combination with ipilimumab:</u> • Administer 1 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then 3 mg/kg intravenously every 2 weeks, or 240 mg every 2 weeks, or 480 mg every 4 weeks until disease progression or unacceptable toxicity
Merkel Cell Carcinoma	<u>Neoadjuvant treatment:</u> • Administer 240 mg intravenously every 2 weeks (days 1 and 15) for a total of 2 doses <u>All other settings:</u> Single agent: • Administer 240 mg intravenously every 2 weeks or 3 mg/kg intravenously every 2 weeks until disease progression or unacceptable toxicity In combination with ipilimumab:

	- Administer 1 mg/kg intravenously OR 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then follow with the single agent regimen; OR - Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity
Non-Small Cell Lung Cancer (NSCLC)	<u>Neoadjuvant treatment followed by optional adjuvant treatment:</u> - Administer 360 mg intravenously with platinum-doublet chemotherapy every 3 weeks for up to 4 cycles with the option of continuing single-agent nivolumab as adjuvant treatment after surgery at 480 mg intravenously every 4 weeks for up to 13 cycles or until disease recurrence or unacceptable toxicity <u>Single agent:</u> - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>In combination with ipilimumab:</u> - Administer 360 mg intravenously every 3 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity for up to 2 years; OR - Administer 3 mg/kg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity for up to 2 years <u>In combination with ipilimumab and platinum-doublet chemotherapy:</u> - Administer 360 mg intravenously every 3 weeks (given in combination with ipilimumab every 6 weeks and histology-based platinum-doublet chemotherapy every 3 weeks for 2 cycles) until disease progression or unacceptable toxicity for up to 2 years.
Pediatric Primary Mediastinal Large B-Cell Lymphoma (PMBCL)	<u>In combination with brentuximab vedotin:</u> Administer 3 mg/kg intravenously, with brentuximab vedotin on day 1, every 2 weeks until disease progression or unacceptable toxicity
Small Bowel Adenocarcinoma	<u>Single agent:</u> - Administer 3 mg/kg intravenously every 2 weeks, or 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks, or 6 mg/kg intravenously every 4 weeks until disease progression or unacceptable toxicity <u>In combination with ipilimumab:</u> - Administer 3 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab on the same day), then 3 mg/kg or 240 mg intravenously every 2 weeks until disease progression or unacceptable toxicity
Small Cell Lung Cancer (SCLC)	Administer 3 mg/kg intravenously every 2 weeks or 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity
Soft Tissue Sarcoma	<u>In combination with ipilimumab:</u>

	Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity
Uterine Neoplasms	- Administer 3 mg/kg intravenously every 2 weeks for 8 doses, then 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity; OR - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity
Vulvar Cancer, Vaginal Cancer & Cervical Cancer	<u>Single agent:</u> - Administer 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity for up to 2 years <u>In combination with ipilimumab:</u> - Administer nivolumab 3 mg/kg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression, unacceptable toxicity or for a maximum of 2 years; OR - Administer nivolumab 1 mg/kg intravenously every 3 weeks for 4 doses (given in combination with ipilimumab every 3 weeks), then followed by single agent regimen until disease progression, unacceptable toxicity or for a maximum of 2 years
CLL/SLL	<u>In combination with ibrutinib:</u> Administer 3 mg/kg intravenously every 2 weeks or 240 mg intravenously every 2 weeks or 480 mg intravenously every 4 weeks, until disease progression or unacceptable toxicity
Squamous Cell Skin Cancer	<u>Weight-based dosing:</u> Administer 3 mg/kg intravenously every 2 weeks until disease progression or unacceptable toxicity for up to 1 year OR <u>Flat dosing:</u> Administer 240mg intravenously every 2 weeks until disease progression or unacceptable toxicity for up to 2 years
Ovarian, Fallopian Tube, & Primary Peritoneal Cancer	<u>Clear cell carcinoma of the ovary:</u> - Administer 3 mg/kg intravenously every 3 weeks for a total of 4 doses (given in combination with ipilimumab on the same day), then 480 mg intravenously every 4 weeks until disease progression or unacceptable toxicity for up to 2 years <u>Small cell carcinoma of the ovary (hypercalcemic type):</u> - Administer 240 mg intravenously every 2 weeks (given in combination with ipilimumab every 6 weeks) until disease progression or unacceptable toxicity

Dosing should be calculated using actual body weight and not flat dosing (as applicable) based on the following:

Frequency (days)	Dosing (mg/kg)	Weight (kg)	Dose (mg)
14	3	<80	220
		<73	200
		<66	180
		<58	160
		<51	140
		<44	120
21	4.5	<80	340
		<78	320
		<73	300
		<68	280
		<63	260
		<58	240
		<53	220
		<48	200
		<44	180
28	6	<80	440
		<77	420
		<73	400
		<69	380
		<66	360
		<62	340
		<58	320
		<55	300
		<51	280
		<47	260
		<44	240

Note: This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide. Member-specific variables should be taken into account.

VI. Billing Code/Availability Information

HCPSC Code:

- J9299 – Injection, nivolumab, 1 mg; 1 billable unit = 1 mg

NDC(s):

- Opdivo 40 mg/4 mL single-dose vial: 00003-3772-xx
- Opdivo 100 mg/10 mL single-dose vial: 00003-3774-xx
- Opdivo 120 mg/12 mL single-dose vial: 00003-3756-xx
- Opdivo 240 mg/24 mL single-dose vial: 00003-3734-xx

VII. References (STANDARD)

1. Opdivo [package insert]. Princeton, NJ; Bristol-Myers Squibb Company; March 2026. Accessed April 2026.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) nivolumab. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
3. Scherpereel A, Mazieres J, Greillier L, et al. Second- or third-line nivolumab (Nivo) versus nivo plus ipilimumab (Ipi) in malignant pleural mesothelioma (MPM) patients: Results of the IFCT-1501 MAPS2 randomized phase II trial. [Abstract]. J Clin Oncol 2017;35: Abstract LBA 8507.
4. Walocko FM, Scheier BY, Harms PW, et al. Metastatic Merkel cell carcinoma response to nivolumab. J Immunother Cancer. 2016 Nov 15;4:79.
5. Tawbi HA-H, Forsyth PAJ, Algazi AP, et al. Combined Nivolumab and Ipilimumab in Melanoma Metastatic to the Brain. N Engl J Med. 2018 Aug 23;379(8):722-730.
6. Morris VK, Salem ME, Nimeiri H, et al. Nivolumab for previously treated unresectable metastatic anal cancer (NCI9673): a multicentre, single-arm, phase 2 study. Lancet Oncol. 2017 Apr;18(4):446-453. doi: 10.1016/S1470-2045(17)30104-3. Epub 2017 Feb 18.
7. Zhao X, Ivaturi V, Gopalakrishnan M, et al. Abstract CT 101: A model-based exposure-response (E-R) assessment of a nivolumab (NIVO) 4-weekly (Q4W) dosing schedule across multiple tumor types. Cancer Res July 1 2017 (77) (13 Supplement) CT101; DOI: 10.1158/1538-7445.AM2017-CT101.
8. Zhao X, Suryawanshi M, Hruska M, et al. Assessment of nivolumab benefit-risk profile of a 240 mg flat dose relative to a 3 mg/kg dosing regimen in patients with advanced tumors. Ann Oncol 2017; 28:2002-2008.

9. Feng Y, Xiaoning W, Bajaj G, et al. Nivolumab exposure-response analyses of efficacy and safety in previously treated squamous or nonsquamous non-small cell lung cancer. *ClinCa Res* 2017;23(18): 5394-5405.
10. Gupta S, Bellmunt J, Plimack ER, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol*. 2022 June 1;40(16_suppl):4577.
11. Hellmann MD, Ciuleanu TE, Pluzanski A, et al. Nivolumab plus ipilimumab in lung cancer with a high tumor mutational burden. *N Engl J Med* 2018; 378:2093-2104.
12. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. *J Oncol Pract*. 2018 Mar;14(3):e130-e136.
13. Hematology/Oncology Pharmacy Association (2019). Intravenous Cancer Drug Waste Issue Brief. Updated February 2024. Available at https://www.hoparx.org/documents/287/HOPA_Drug_Waste_Issue_Brief_-_Updated_02.22.24.
14. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. *BMJ*. 2016 Feb 29;352:i788.
15. Weber JS, D'Angelo SP, Minor D, et al. Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol*. 2015 Apr;16(4):375-84. doi: 10.1016/S1470-2045(15)70076-8. Epub 2015 Mar 18.
16. Robert C, Long GV, Brady B, et al. Nivolumab in previously untreated melanoma without BRAF mutation. *N Engl J Med*. 2015 Jan 22;372(4):320-30. doi: 10.1056/NEJMoa1412082. Epub 2014 Nov 16.
17. Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N Engl J Med*. 2015 Jul 2;373(1):23-34. doi: 10.1056/NEJMoa1504030. Epub 2015 May 31.
18. Weber J, Mandala M, Del Vecchio M, et al. Adjuvant Nivolumab versus Ipilimumab in Resected Stage III or IV Melanoma. *N Engl J Med*. 2017 Nov 9;377(19):1824-1835. doi: 10.1056/NEJMoa1709030. Epub 2017 Sep 10.
19. Algazi AP, Tsai KK, Shoushtari AN, et al. Clinical outcomes in metastatic uveal melanoma treated with PD-1 and PD-L1 antibodies. *Cancer*. 2016 Nov 15;122(21):3344-3353. doi: 10.1002/cncr.30258. Epub 2016 Aug 17.
20. Piulats JM, Espinosa E, de la Cruz-Merino L, et al. Nivolumab plus ipilimumab for treatment-naive metastatic uveal melanoma: An open-label, multicenter, phase II trial by the Spanish Multidisciplinary Melanoma Group (GEM-1402). *J Clin Oncol* 2021; 39:586-598.c
21. El-Khoueiry AB, Sangro B, Yau T, et al. Nivolumab in patients with advanced hepatocellular carcinoma (CheckMate 040): an open-label, non-comparative, phase 1/2 dose escalation and expansion trial. *Lancet*. 2017 Jun 24;389(10088):2492-2502. doi: 10.1016/S0140-6736(17)31046-2. Epub 2017 Apr 20.

22. Brahmer J, Reckamp KL, Baas P, et al. Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer. *N Engl J Med*. 2015 Jul 9;373(2):123-35. doi: 10.1056/NEJMoa1504627. Epub 2015 May 31.
23. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med*. 2015 Oct 22;373(17):1627-39. doi: 10.1056/NEJMoa1507643. Epub 2015 Sep 27.
24. Antonia SJ, López-Martin JA, Bendell J, et al. Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase 1/2 trial. *Lancet Oncol*. 2016 Jul;17(7):883-895. doi: 10.1016/S1470-2045(16)30098-5. Epub 2016 Jun 4.
25. Motzer RJ, Escudier B, McDermott DF, et al. Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2015 Nov 5;373(19):1803-13. doi: 10.1056/NEJMoa1510665. Epub 2015 Sep 25.
26. Motzer RJ, Tannir NM, McDermott DF, et al. Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2018 Apr 5;378(14):1277-1290. doi: 10.1056/NEJMoa1712126. Epub 2018 Mar 21.
27. Armand P, Engert A, Younes A, et al. Nivolumab for Relapsed/Refractory Classic Hodgkin Lymphoma After Failure of Autologous Hematopoietic Cell Transplantation: Extended Follow-Up of the Multicohort Single-Arm Phase II CheckMate 205 Trial. *J Clin Oncol*. 2018 May 10;36(14):1428-1439. doi: 10.1200/JCO.2017.76.0793. Epub 2018 Mar 27.
28. Ansell SM, Lesokhin AM, Borrello I, et al. PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma. *N Engl J Med*. 2015 Jan 22;372(4):311-9. doi: 10.1056/NEJMoa1411087. Epub 2014 Dec 6.
29. Ferris RL, Blumenschein G Jr, Fayette J, et al. Nivolumab for Recurrent Squamous-Cell Carcinoma of the Head and Neck. *N Engl J Med*. 2016 Nov 10;375(19):1856-1867. Epub 2016 Oct 8.
30. Sharma P, Retz M, Siefker-Radtke A, et al. Nivolumab in metastatic urothelial carcinoma after platinum therapy (CheckMate 275): a multicentre, single-arm, phase 2 trial. *Lancet Oncol*. 2017 Mar;18(3):312-322. doi: 10.1016/S1470-2045(17)30065-7. Epub 2017 Jan 26.
31. Overman MJ, McDermott R, Leach JL, et al. Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study. *Lancet Oncol*. 2017 Sep;18(9):1182-1191. doi: 10.1016/S1470-2045(17)30422-9. Epub 2017 Jul 19.
32. Overman MJ, Lonardi S, Wong KYM, et al. Durable Clinical Benefit With Nivolumab Plus Ipilimumab in DNA Mismatch Repair-Deficient/Microsatellite Instability-High Metastatic Colorectal Cancer. *J Clin Oncol*. 2018 Mar 10;36(8):773-779. doi: 10.1200/JCO.2017.76.9901. Epub 2018 Jan 20.
33. Topalian SL, Bhatia S, Hollebecque A, et al. Non-comparative, open-label, multiple cohort, phase 1/2 study to evaluate nivolumab (NIVO) in patients with virus-associated tumors

- (CheckMate 358): Efficacy and safety in Merkel cell carcinoma (MCC). DOI: 10.1158/1538-7445.AM2017-CT074 Published July 2017.
34. Long GV, Atkinson V, Lo S, et al. Combination nivolumab and ipilimumab or nivolumab alone in melanoma brain metastases: a multicentre randomised phase 2 study. *Lancet Oncol*. 2018 May;19(5):672-681. doi: 10.1016/S1470-2045(18)30139-6. Epub 2018 Mar 27.
35. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Anal Carcinoma. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
36. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Gestational Trophoblastic Neoplasia. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
37. Scherpereel A, Mazieres J, Greillier L, et al. Nivolumab or nivolumab plus ipilimumab in patients with relapsed malignant pleural mesothelioma (IFCT-1501 MAPS2): a multicentre, open-label, randomised, non-comparative, phase 2 trial. *Lancet Oncol*. 2019 Feb;20(2):239-253. doi: 10.1016/S1470-2045(18)30765-4. Epub 2019 Jan 16.
38. Disselhorst MJ, Quispel-Janssen J, Lalezari F, et al. Ipilimumab and nivolumab in the treatment of recurrent malignant pleural mesothelioma (INITIATE): results of a prospective, single-arm, phase 2 trial. *Lancet Respir Med*. 2019 Mar;7(3):260-270. doi: 10.1016/S2213-2600(18)30420-X. Epub 2019 Jan 16.
39. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Small Bowel Adenocarcinoma. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
40. Chan TSY, Li J, Loong F, et al. PD1 blockade with low-dose nivolumab in NK/T cell lymphoma failing L-asparaginase: efficacy and safety. *Ann Hematol*. 2018 Jan;97(1):193-196. doi: 10.1007/s00277-017-3127-2. Epub 2017 Sep 6.

41. Goldman JW, Crino L, Vokes EE, et al. P2.36 Nivolumab (nivo) in patients (pts) with advanced (adv) NSCLC and central nervous system (CNS) metastases (mets). *J Thorac Oncol* 2016;11:S238-S239.
42. Gauvain C, Vauleon E, Chouaid C, et al. Intracerebral efficacy and tolerance of nivolumab in non–small-cell lung cancer patients with brain metastases. *Lung Cancer*. 2018 Feb; 116:62-66. doi: 10.1016/j.lungcan.2017.12.008.
43. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Non-Small Cell Lung Cancer. Version 5.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
44. Kato K, Cho BC, Takahashi M, et al. Nivolumab versus chemotherapy in patients with advanced esophageal squamous cell carcinoma refractory or intolerant to previous chemotherapy (ATTRACTION-3): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol*. 2019;20(11):1506-1517. doi:10.1016/S1470-2045(19)30626-6.
45. Hellmann MD, Paz-Ares L, Bernabe Caro R, et al. Nivolumab plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer. *N Engl J Med*. 2019;381(21):2020-2031. doi:10.1056/NEJMoa1910231.
46. Reck M, Ciuleanu T-E, Dols MC, et al. Nivolumab (NIVO) + ipilimumab (IPI) + 2 cycles of platinum-doublet chemotherapy (chemo) vs 4 cycles chemo as first-line (1L) treatment (tx) for stage IV/recurrent non-small cell lung cancer (NSCLC): CheckMate 9LA [abstract]. *J Clin Oncol* 2020;38:Abstract 9501-9501.
47. Zalcman G, Peters S, Mansfield AS, et al. Checkmate 743: A phase 3, randomized, open-label trial of nivolumab (nivo) plus ipilimumab (ipi) vs pemetrexed plus cisplatin or carboplatin as first-line therapy in unresectable pleural mesothelioma. *Journal of Clinical Oncology* 2017 35:15_suppl, TPS8581-TPS8581.
48. Azad NS, Gray RJ, Overman MJ, et al. Nivolumab Is Effective in Mismatch Repair-Deficient Noncolorectal Cancers: Results From Arm Z1D-A Subprotocol of the NCI-MATCH (EAY131) Study. *J Clin Oncol*. 2020 Jan 20;38(3):214-222.
49. Naumann RW, Hollebecque A, Meyer T, et al. Safety and Efficacy of Nivolumab Monotherapy in Recurrent or Metastatic Cervical, Vaginal, or Vulvar Carcinoma: Results From the Phase I/II CheckMate 358 Trial. *J Clin Oncol*. 2019 Nov 1;37(31):2825-2834.
50. Choueiri TK, Powles T, Burotto M, et al. 696O_PR Nivolumab + cabozantinib vs sunitinib in first-line treatment for advanced renal cell carcinoma: First results from the randomized phase III CheckMate 9ER trial. Volume 31, SUPPLEMENT 4, S1159, September 01, 2020.

51. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Bladder Cancer. Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
52. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Esophageal and Esophagogastric Junction Cancers. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
53. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Gastric Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
54. Herrera AF, Moskowitz AJ, Bartlett NL, et al. Interim results of brentuximab vedotin in combination with nivolumab in patients with relapsed or refractor Hodgkin lymphoma. *Blood*. 2018 Mar 15;131 (11):1183-1194.
55. Cole PD, Mauz-Körholz C, Mascarini M, et al. HL-032: Nivolumab and Brentuximab Vedotin (BV)–Based, Response-Adapted Treatment in Children, Adolescents, and Young Adults (CAYA) With Standard-Risk Relapsed/Refractory Classical Hodgkin Lymphoma (R/R cHL): Primary Analysis of the Standard-Risk Cohort of the Phase 2 CheckMate 744 Study. *Clinical Lymphoma Myeloma and Leukemia*. Volume 20, Supplement 1, September 2020, Pages S245-S246.
56. Moehler M, Shitara K, Garrido M, et al. Nivolumab (nivo) plus chemotherapy (chemo) versus chemo as first-line (1L) treatment for advanced gastric cancer/gastroesophageal junction cancer (GC/GEJC)/esophageal adenocarcinoma (EAC): First results of the CheckMate 649 study. [abstract]. Presented at the Oral Presentation presented at the ESMO 2020 Annual Meeting; September 19-21, 2020; Virtual Meeting.
57. Kelly RJ, Ajani JA, Kuzdzal J, et al. Adjuvant Nivolumab in Resected Esophageal or Gastroesophageal Junction Cancer. *N Engl J Med*. 2021 Apr 1;384(13):1191-1203. doi: 10.1056/NEJMoa2032125.
58. Nivolumab. Micromedex Solutions. Greenwood Village, CO: Truven Health Analytics. <http://micromedex.com/>. Updated May 29, 2025. Accessed May 2025.

59. Lenz HJ, Lonardi S, Zagonel V, et al. Nivolumab (NIVO) + low-dose ipilimumab (IPI) as first-line (1L) therapy in microsatellite instability-high/DNA mismatch repair deficient (MSI-H/dMMR) metastatic colorectal cancer (mCRC): Clinical update [abstract]. *Journal of Clinical Oncology* 2019;37:3521-3521.
60. Bellmunt, J. (2025). Treatment of metastatic urothelial cancer of the bladder and urinary tract. In Lerner SP, Shah S (Eds.), *UptoDate*. Last updated Feb 05, 2026. Accessed April 13, 2026. Available from <https://www.uptodate.com/contents/treatment-of-metastatic-urothelial-cancer-of-the-bladder-and-urinary-tract>.
61. Ready NE, Ott PA, Hellmann MD, et al. Nivolumab Monotherapy and Nivolumab Plus Ipilimumab in Recurrent Small Cell Lung Cancer: Results From the CheckMate 032 Randomized Cohort. *J Thorac Oncol*. 2020 Mar;15(3):426-435. doi: 10.1016/j.jtho.2019.10.004.
62. Bajorin DF, Witjes JA, Gschwend JE, et al. Adjuvant Nivolumab versus Placebo in Muscle-Invasive Urothelial Carcinoma. *N Engl J Med*. 2021 Jun 3;384(22):2102-2114. doi: 10.1056/NEJMoa2034442.
63. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Cervical Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
64. Fennell DA, Ewings S, Ottensmeier C, et al. Nivolumab versus placebo in patients with relapsed malignant mesothelioma (CONFIRM): a multicentre, double-blind, randomised, phase 3 trial. *Lancet Oncol* 2021; 22:1530.
65. Topalian SL, Bhatia S, Amin A, et al. Neoadjuvant Nivolumab for Patients With Resectable Merkel Cell Carcinoma in the CheckMate 358 Trial. *J Clin Oncol*. 2020;38(22):2476-2487. doi:10.1200/JCO.20.00201.
66. Forde PM, Spicer J, Lu S et al. Neoadjuvant Nivolumab plus Chemotherapy in Resectable Lung Cancer. *N Engl J Med*. 2022 May 26;386(21):1973-1985. doi: 10.1056/NEJMoa2202170. Epub 2022 Apr 11. PMID: 35403841; PMCID: PMC9844511
67. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Mesothelioma: Peritoneal. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
68. Scherpereel A, Mazieres J, Greillier L, et al; French Cooperative Thoracic Intergroup. Nivolumab or nivolumab plus ipilimumab in patients with relapsed malignant pleural mesothelioma (IFCT-1501 MAPS2): a multicentre, open-label, randomised, non-comparative,

- phase 2 trial. *Lancet Oncol.* 2019 Feb;20(2):239-253. doi: 10.1016/S1470-2045(18)30765-4. Epub 2019 Jan 16. Erratum in: *Lancet Oncol.* 2019 Mar;20(3):e132.
69. Doki Y, Ajani JA, Kato K, et al. Nivolumab Combination Therapy in Advanced Esophageal Squamous-Cell Carcinoma. *N Engl J Med.* 2022 Feb 3;386(5):449-462. doi: 10.1056/NEJMoa2111380.
 70. De Santis M, Bellmunt J, Mead G, et al. Randomized phase II/III trial assessing gemcitabine/carboplatin and methotrexate/carboplatin/vinblastine in patients with advanced urothelial cancer “unfit” for cisplatin-based chemotherapy: phase II—results of EORTC study 30986. *J Clin Oncol.* 2009 Nov 20;27(33):5634-9. Doi: 10.1200/JCO.2008.21.4924. Epub 2009 Sep 28.
 71. Bouffet E, Larouche V, Campbell BB, et al. Immune Checkpoint Inhibition for Hypermutant Glioblastoma Multiforme Resulting From Germline Biallelic Mismatch Repair Deficiency. *J Clin Oncol.* 2016 Jul 1;34(19):2206-11.
 72. Schenker M, Burotto M, Richardet M, et al. Randomized, open-label, phase 2 study of nivolumab plus ipilimumab or nivolumab monotherapy in patients with advanced or metastatic solid tumors of high tumor mutational burden. *J Immunother Cancer* 2024; 12:e008872
 73. MG, Lee HJ, Palmer J, et al. Response-adapted anti-PD-1-based salvage therapy for Hodgkin lymphoma with nivolumab alone or in combination with ICE. *Blood.* 2022 Jun 23;139(25):3605-3616. doi: 10.1182/blood.2022015423.
 74. Zinzani P, Santoro A, Gritti G, et al. Nivolumab Combined With Brentuximab Vedotin for Relapsed/Refractory Primary Mediastinal Large B-Cell Lymphoma: Efficacy and Safety From the Phase II CheckMate 436 Study. *J Clin Oncol.* 2019 Nov 20;37(33):3081-3089. doi: 10.1200/JCO.19.01492. Epub 2019 Aug 9.
 75. Davis K, Fox E, Merchant M, et al. Nivolumab in children and young adults with relapsed or refractory solid tumours or lymphoma (ADVIL1412): a multicentre, open-label, single-arm, phase 1–2 trial. *The Lancet.* volume 21, issue 4, p541-550, April 01, 2020 [https://doi.org/10.1016/S1470-2045\(20\)30023-1](https://doi.org/10.1016/S1470-2045(20)30023-1).
 76. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Pediatric Aggressive Mature B-Cell Lymphomas. Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
 77. Younes A, Santoro A, Shipp M, et al. Nivolumab for classical Hodgkin’s lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: a multicentre, multicohort, single-arm phase 2 trial. *Lancet Oncol.* 2016 Sep;17(9):1283-94. doi: 10.1016/S1470-2045(16)30167-X.
 78. Chung C, Li J, Steuer C, et al. Phase II Multi-institutional Clinical Trial Result of Concurrent Cetuximab and Nivolumab in Recurrent and/or Metastatic Head and Neck Squamous Cell

- Carcinoma. Clin Cancer Res. 2022 Jun 1;28(11):2329-2338. doi: 10.1158/1078-0432.CCR-21-3849.
79. Zer A, Icht O, Yosef L, et al. Phase II single-arm study of nivolumab and ipilimumab (Nivo/Ipi) in previously treated classical Kaposi sarcoma (cKS). Annals of Oncology. Volume 33, Issue 7, July 2022, Pages 720-727. <https://doi.org/10.1016/j.annonc.2022.03.012>.
 80. Pelster MS, Gruschkus SK, Bassett R, et al. Nivolumab and Ipilimumab in Metastatic Uveal Melanoma: Results From a Single-Arm Phase II Study. J Clin Oncol. 2021 Feb 20;39(6):599-607. doi: 10.1200/JCO.20.00605.
 81. Baas P, Scherpereel A, Nowak AK, et al. First-line nivolumab plus ipilimumab in unresectable malignant pleural mesothelioma (CheckMate 743): a multicentre, randomised, open-label, phase 3 trial. Lancet. 2021 Jan 30;397(10272):375-386. doi: 10.1016/S0140-6736(20)32714-8.
 82. Blank CU, Rozeman EA, Fanchi LF, et al. Neoadjuvant versus adjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma. Nat Med. 2018 Nov;24(11):1655-1661. doi: 10.1038/s41591-018-0198-0.
 83. Glutsch V, Kneitz, Gesierich A, et al. Activity of ipilimumab plus nivolumab in avelumab-refractory Merkel cell carcinoma. Cancer Immunology, Immunotherapy volume 70, pages2087–2093 (2021)
 84. Wagner M, Othus M, Patel S, et al. Multicenter phase II trial (SWOG S1609, cohort 51) of ipilimumab and nivolumab in metastatic or unresectable angiosarcoma: a substudy of dual anti-CTLA-4 and anti-PD-1 blockade in rare tumors (DART). J Immunother Cancer. 2021 Aug;9(8):e002990. doi: 10.1136/jitc-2021-002990.
 85. Kim S, Wuthrick E, Blakaj D, et al. Combined nivolumab and ipilimumab with or without stereotactic body radiation therapy for advanced Merkel cell carcinoma: a randomized, open label, phase 2 trial. The Lancet. Published: September 11, 2022. doi:[https://doi.org/10.1016/S0140-6736\(22\)01659-2](https://doi.org/10.1016/S0140-6736(22)01659-2). PlumX Metrics
 86. Yau T, Park JW, Finn RS, et al. Nivolumab versus sorafenib in advanced hepatocellular carcinoma (CheckMate 459): a randomised, multicentre, open-label, phase 3 trial. Lancet Oncol. 2022 Jan;23(1):77-90.
 87. Kudo M, Matilla A, Santoro A, et al. CheckMate 040 cohort 5: A phase I/II study of nivolumab in patients with advanced hepatocellular carcinoma and Child-Pugh B cirrhosis. J Hepatol. 2021 Sep;75(3):600-609.
 88. Long GV, Del Vecchio M, Weber J, et al. (2023). Adjuvant therapy with nivolumab versus placebo in patients with resected stage IIB/C melanoma (CheckMate 76K). SKIN The Journal of Cutaneous Medicine, 7(2), s163. <https://doi.org/10.25251/skin.7.suppl.163>.
 89. Advani RH, Moskowitz AJ, Bartlett NL, et al. Brentuximab vedotin in combination with nivolumab in relapsed or refractory Hodgkin lymphoma: 3-year study results. Blood. 2021 Aug 12;138(6):427-438. Doi: 10.1182/blood.2020009178.

90. Dagogo-Jack I, Madison RW, Lennerz JK, et al. Molecular characterization of mesothelioma: Impact of histologic type and site of origin on molecular landscape. *JCO Precis Oncol* 2022;6:e2100422.
91. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Colon Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
92. van der Heijden, MS, Sonpavde G, Powles T, et al; CheckMate 901 Trial Investigators. Nivolumab plus Gemcitabine-Cisplatin in Advanced Urothelial Carcinoma. *N Engl J Med*. 2023 Nov 9;389(19):1778-1789. doi: 10.1056/NEJMoa2309863. Epub 2023 Oct 22. PMID: 37870949.
93. Amaria R, Reddy S, Tawbi H, et al. Neoadjuvant Immune Checkpoint Blockade in High-Risk Resectable Melanoma. *Nat Med*. 2018 Nov; 24(11): 1649–1654. Published online 2018 Oct 8. Doi: 10.1038/s41591-018-0197-1
94. Ma, D, Ding X, Zhang C, et al Combined targeted therapy and immunotherapy in anaplastic thyroid carcinoma with distant metastasis: A case report. *World J Clin Cases* 2022;10:3849-3855.
95. Kollipara R, Schneider K, Radovich M, et al. Exceptional response with immunotherapy in a patient with anaplastic thyroid cancer. *Oncologist* 2017;22:1149-1151.
96. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Thyroid Carcinoma. Version 1.2025. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
97. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Vaginal Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
98. Reijers ILM, Menzies AM, van Akkooi ACJ, et al. Personalized response-directed surgery and adjuvant therapy after neoadjuvant ipilimumab and nivolumab in high-risk stage III melanoma: the PRADO trial. *Nat Med* 2022;28:1178-1188.

99. Versluis JM, Menzies AM, Sikorska K, et al. Survival update of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma in the OpACIN and OpACINneo trials. *Ann Oncol* 2023;34:420-430.
100. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Melanoma: Cutaneous Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
101. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab followed by Nivolumab: Colon Cancer Chemotherapy Order Template, COL68. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
102. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab followed by Nivolumab: Rectal Cancer Chemotherapy Order Template, REC80. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
103. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for mFOLFOX6 (Continuous Infusion Fluorouracil/Leucovorin/OXALlplatin) + Nivolumab: Gastric Cancer Chemotherapy Order Template, GAS95. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
104. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for CapeOX (Capecitabine/OXALlplatin) + Nivolumab: Gastric Cancer Chemotherapy Order Template, GAS96. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
105. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab followed by nivolumab: Ampullary Adenocarcinoma Chemotherapy Order Template, AMP22. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES®

are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.

106. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Mesothelioma: Peritoneal Chemotherapy Order Template, MPEM10. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
107. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab followed by Nivolumab: Central Nervous System Cancers Chemotherapy Order Template, CNS61. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
108. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Central Nervous System Cancers Chemotherapy Order Template, CNS63. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
109. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab: Non-Small Cell Lung Cancer Chemotherapy Order Template, NSC97. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
110. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: T-Cell Lymphomas Chemotherapy Order Template, TCL39. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
111. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Small Cell Lung Cancer Chemotherapy Order Template, SCL24. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National

Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.

112. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Endometrial Carcinoma Chemotherapy Order Template, UTE34. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
113. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Cervical Cancer Chemotherapy Order Template, CRV35. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
114. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Vulvar Cancer Chemotherapy Order Template, VUL17. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
115. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Vaginal Cancer Chemotherapy Order Template, VAG34. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
116. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Brentuximab vedotin + Nivolumab: Hodgkin Lymphoma Chemotherapy Order Template, HDL53. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
117. Herrera AF, LeBlanc ML, Castellino SM, et al. Nivolumab(N)+AVD in advanced stage (AS) classic Hodgkin's lymphoma (HL). *N Engl J Med* 2024;391:1379-1389.
118. Bröckelmann PJ, Buhen I, Meissner J et al. Nivolumab and Doxorubicin, Vinblastine, and Dacarbazine in Early-Stage Unfavorable Hodgkin Lymphoma: Final Analysis of the Randomized German Hodgkin Study Group Phase II NIVAHG Trial. *J Clin Oncol.* 2023 Feb 20;41(6):1193-1199. doi: 10.1200/JCO.22.02355. Epub 2022 Dec 12. PMID: 36508302.

119. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for AVD (DOXOrubicin/VinBLAStine/Dacarbazine) + Nivolumab: Hodgkin Lymphoma Chemotherapy Order Template, HDL75. National Comprehensive Cancer Network, 2025. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2025.
120. Cascone T, Awad MM, Spicer JD, et al. Perioperative Nivolumab in Resectable Lung Cancer. *N Engl J Med*. 2024 May 16;390(19):1756-1769. doi: 10.1056/NEJMoa2311926. PMID: 38749033.
121. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Kaposi Sarcoma Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
122. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2026.
123. Younes A, Brody J, Carpio C, et al. Safety and activity of ibrutinib in combination with nivolumab in patients with relapsed non-Hodgkin lymphoma or chronic lymphocytic leukaemia: a phase 1/2a study. *Lancet Haematol* 2019;6:e67-e78.
124. Lebbe C, Meyer N, Mortier L, et al. Initial results from a phase IIIb/IV study evaluating two dosing regimens of nivolumab (NIVO) in combination with ipilimumab (IPI) in patients with advanced melanoma (CheckMate 511)(abstract). *Ann Oncol* 2018;29:LBA47.
125. Andre T, Elez E, Van Cutsem E, et al; CheckMate 8HW Investigators. Nivolumab plus ipilimumab in Microsatellite-Instability-High Metastatic Colorectal Cancer. *N Engl J Med*. 2024 Nov 28;391(21):2014-2026. doi: 10.1056/NEJMoa2402141. PMID: 39602630.
126. Yau T, Galle PR, Decaens T, et al. Nivolumab plus ipilimumab versus lenvatinib or sorafenib as first-line treatment for unresectable hepatocellular carcinoma (CheckMate 9DW): An open-label, randomised, phase 3 trial. *Lancet* 2025;405:1851-1864.
127. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Ipilimumab + Nivolumab: Hepatocellular Carcinoma Chemotherapy Order Template, HEP33. National Comprehensive Cancer Network, 2025. NATIONAL

COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2025.

128. Patel SP, Othus M, Chae YK, et al. Phase II basket trial of dual anti-CTLA-4 and anti-PD-1 blockade in rare tumors (DART) SWOG S1609: Adrenocortical carcinoma cohort. *J Immunother Cancer* 2024;12:e009074.
129. Munhoz RR, Nader-Marta G, de Camargo VP, et al. A phase 2 study of first-line nivolumab in patients with locally advanced or metastatic cutaneous squamous-cell carcinoma. *Cancer*. 2022;128(24):4223-4231. doi:10.1002/cncr.34463
130. Lang R, Welpner T, Richtig E, et al. Nivolumab for locally advanced and metastatic cutaneous squamous cell carcinoma (NIVOSQUACS study)-Phase II data covering impact of concomitant haematological malignancies. *J Eur Acad Dermatol Venereol*. 2023;37(9):1799-1810. doi:10.1111/jdv.19218
131. Harker-Murray P, Mauz-Korholz C, Leblanc T, et al. Nivolumab and brentuximab vedotin with or without bendamustine for R/R Hodgkin lymphoma in children, adolescents, and young adults. *Blood* 2023;141:2075-2084.
132. Greve P, Beishuizen A, Hagleitner M, et al. Nivolumab plus brentuximab vedotin +/- bendamustine combination therapy: A safe and effective treatment in pediatric recurrent and refractory classical Hodgkin lymphoma. *Front Immunol* 2023;14:1229558.
133. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Appendiceal Neoplasms and Cancers. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
134. Andre T, Tougeron D, Piessen G, et al. Neoadjuvant nivolumab plus ipilimumab and adjuvant nivolumab in localized deficient mismatch repair/microsatellite instability-high gastric or esophagogastric junction adenocarcinoma: The GERCOR NEONIPIGA phase II study. *J Clin Oncol* 2023;41:255-265.
135. Janjigian YY, Shitara K, Moehler M, et al. First-line nivolumab plus chemotherapy versus chemotherapy alone for advanced gastric, gastro-oesophageal junction, and oesophageal adenocarcinoma (CheckMate 649): a randomised, open-label, phase 3 trial. *Lancet* 2021;398:27-40.
136. Oaknin A, Moore K, Meyer T, et al. Nivolumab with or without ipilimumab in patients with recurrent or metastatic cervical cancer (CheckMate 358): a phase 1-2, open-label, multicohort trial. *Lancet Oncol* 2024;25:588-602.

137. Palmerini E, Lopez Pousa A, Grignani G, et al. Nivolumab and sunitinib in patients with advanced bone sarcomas: a multicenter, single-arm, phase 2 trial. *Cancer* 2025;303:e35628.
138. Long GV, Tykodi SS, Schneider JG, et al. Assessment of nivolumab exposure and clinical safety of 480 mg every 4 weeks flat-dosing schedule in patients with cancer. *Ann Oncol.* 2018;29(11):2208-2213.
139. Gao B, Carlino MS, Michael M, et al. Nivolumab and Ipilimumab Combination Treatment in Advanced Ovarian and Endometrial Clear Cell Cancers: A Nonrandomized Clinical Trial. *JAMA Oncol.* 2025 Sep 1;11(9):982-989.
140. Chae YK, Othus M, Patel, SP, et al. A phase II basket trial of dual anti-CTLA-4 and anti-PD-1 blockade in rare tumors (DART) SWOG S1609: durable responses and delayed pseudoprogression in small cell carcinoma of the ovary, hypercalcemic type cohort. *Cancer Commun (Lond).* 2025 Aug;45(8):976-986.
141. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Ovarian Cancer Including Fallopian Tube Cancer and Primary Peritoneal Cancer. Version 4.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.

VIII. References (ENHANCED)

- 1e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Central Nervous System Cancers. Version 3.2025. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 2e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Head and Neck Cancers. Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 3e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Hepatocellular Carcinoma. Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®.

NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.

- 4e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Hodgkin Lymphoma. Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 5e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Kidney Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 6e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Mesothelioma: Pleural. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 7e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Merkel Cell Carcinoma. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 8e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Rectal Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 9e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Small Cell Lung Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES®

are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.

- 10e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) T-Cell Lymphomas. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 11e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Melanoma: Uveal. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 12e. Robert C, Schachter J, Long GV, et al. Pembrolizumab versus Ipilimumab in Advanced Melanoma. *N Engl J Med*. 2015 Jun 25;372(26):2521-32. Doi: 10.1056/NEJMoa1503093.
- 13e. Ascierto PA, Long GV, Robert C, et al. Survival Outcomes in Patients With Previously Untreated BRAF Wild-Type Advanced Melanoma Treated With Nivolumab Therapy: Three-Year Follow-up of a Randomized Phase 3 Trial [published correction appears in *JAMA Oncol*. 2019 Feb 1;5(2):271]. *JAMA Oncol*. 2019;5(2):187–194. Doi:10.1001/jamaoncol.2018.4514.
- 14e. Wolchok JD, Chiarion-Sileni V, Gonzalez R, et al. Overall Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma [published correction appears in *N Engl J Med*. 2018 Nov 29;379(22):2185]. *N Engl J Med*. 2017;377(14):1345–1356. Doi:10.1056/NEJMoa1709684.
- 15e. Regan MM, Werner L, Rao S, et al. Treatment-Free Survival: A Novel Outcome Measure of the Effects of Immune Checkpoint Inhibition-A Pooled Analysis of Patients With Advanced Melanoma. *J Clin Oncol*. 2019;37(35):3350-3358. Doi:10.1200/JCO.19.00345.
- 16e. Ribas A, Puzanov I, Dummer R, et al. Pembrolizumab versus investigator-choice chemotherapy for ipilimumab-refractory melanoma (KEYNOTE-002): a randomised, controlled, phase 2 trial. *Lancet Oncol*. 2015 Aug;16(8):908-18. Doi: 10.1016/S1470-2045(15)00083-2.
- 17e. Hamid O, Puzanov I, Dummer R, et al. Final analysis of a randomised trial comparing pembrolizumab versus investigator-choice chemotherapy for ipilimumab-refractory advanced melanoma. *Eur J Cancer*. 2017 Nov;86:37-45. Doi: 10.1016/j.ejca.2017.07.022.
- 18e. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma [published correction appears in *N Engl J Med*. 2010 Sep 23;363(13):1290]. *N Engl J Med*. 2010;363(8):711–723. Doi:10.1056/NEJMoa1003466.
- 19e. Larkin J, Minor D, D'Angelo S, et al. Overall Survival in Patients With Advanced Melanoma Who Received Nivolumab Versus Investigator's Choice Chemotherapy in CheckMate 037: A

Randomized, Controlled, Open-Label Phase III Trial. *J Clin Oncol*. 2018;36(4):383–390. Doi:10.1200/JCO.2016.71.8023.

- 20e. Hersh EM, O'Day SJ, Ribas A, et al. A phase 2 clinical trial of nab-paclitaxel in previously treated and chemotherapy-naïve patients with metastatic melanoma. *Cancer*. 2010 Jan 1;116(1):155-63.
- 21e. Kottschade LA, Suman VJ, Amatruda T 3rd, et al. A phase II trial of nab-paclitaxel (ABI-007) and carboplatin in patients with unresectable stage IV melanoma: a North Central Cancer Treatment Group Study, N057E(1). *Cancer*. 2011 Apr 15;117(8):1704-10.
- 22e. Agarwala SS, et al. Randomized phase III study of paclitaxel plus carboplatin with or without sorafenib as second-line treatment in patients with advanced melanoma. *Journal of Clinical Oncology* 2007 25:18_suppl, 8510-8510.
- 23e. Rao RD, et al. Combination of paclitaxel and carboplatin as second-line therapy for patients with metastatic melanoma. *Cancer*. 2006 Jan 15;106(2):375-82.
- 24e. Middleton MR, et al. Randomized phase III study of temozolomide versus dacarbazine in the treatment of patients with advanced metastatic malignant melanoma. *J Clin Oncol*. 2000 Jan;18(1):158-66.
- 25e. Einzig AI, et al. A phase II study of taxol in patients with malignant melanoma. *Invest New Drugs*. 1991 Feb;9(1):59-64.
- 26e. Robert C, Ribas A, Schachter J, et al. Pembrolizumab versus ipilimumab in advanced melanoma (KEYNOTE-006): post-hoc 5-year results from an open-label, multicentre, randomised, controlled, phase 3 study. *Lancet Oncol* 2019; 20:1239.
- 27e. Eggermont AMM, Blank CU, Mandala M, et al. Adjuvant Pembrolizumab versus Placebo in Resected Stage III Melanoma. *N Engl J Med*. 2018 May 10;378(19):1789-1801. doi: 10.1056/NEJMoa1802357.
- 28e. Eggermont AM, Chiarion-Sileni V, Grob JJ, et al. Prolonged Survival in Stage III Melanoma with Ipilimumab Adjuvant Therapy [published correction appears in *N Engl J Med*. 2018 Nov 29;379(22):2185]. *N Engl J Med*. 2016;375(19):1845–1855. doi:10.1056/NEJMoa1611299.
- 29e. Kottschade LA, McWilliams RR, Markovic SN, et al. The use of pembrolizumab for the treatment of metastatic uveal melanoma. *Melanoma Res*. 2016 Jun;26(3):300-3. doi: 10.1097/CMR.0000000000000242.
- 30e. Zimmer L, Vaubel J, Mohr P, et al. Phase II DeCOG-study of ipilimumab in pretreated and treatment-naïve patients with metastatic uveal melanoma. *PLoS One*. 2015;10(3):e0118564. Published 2015 Mar 11. doi:10.1371/journal.pone.0118564.
- 31e. Piulats Rodriguez JM, Ochoa de Olza M, Codes M, et al. Phase II study evaluating ipilimumab as a single agent in the first-line treatment of adult patients (Pts) with metastatic uveal melanoma (MUM): The GEM-1 trial. *J Clin Oncol* 2014; 32S:ASCO #9033.

- 32e. Luke JJ, Callahan MK, Postow MA, et al. Clinical activity of ipilimumab for metastatic uveal melanoma: a retrospective review of the Dana-Farber Cancer Institute, Massachusetts General Hospital, Memorial Sloan-Kettering Cancer Center, and University Hospital of Lausanne experience. *Cancer*. 2013;119(20):3687–3695. doi:10.1002/cncr.28282.
- 33e. Zhu AX, Finn RS, Edeline J, et al. Pembrolizumab in patients with advanced hepatocellular carcinoma previously treated with sorafenib (KEYNOTE-224): a non-randomised, open-label phase 2 trial. *Lancet Oncol*. 2018 Jul;19(7):940-952. doi: 10.1016/S1470-2045(18)30351-6.
- 34e. Crocenzi TS, El-Khoueiry AB, Yau T, et al. Nivolumab (nivo) in sorafenib (sor)-naive and -experienced pts with advanced hepatocellular carcinoma (HCC): CheckMate 040 study. *J Clin Oncol* 35, 2017 (suppl; abstr 4013).
- 35e. Qin S, Bai Y, Lim HY, et al. Randomized, multicenter, open-label study of oxaliplatin plus fluorouracil/leucovorin versus doxorubicin as palliative chemotherapy in patients with advanced hepatocellular carcinoma from Asia. *J Clin Oncol*. 2013 Oct 1;31(28):3501-8. doi: 10.1200/JCO.2012.44.5643.
- 36e. Bruix J, Qin S, Merle P, et al. Regorafenib for patients with hepatocellular carcinoma who progressed on sorafenib treatment (RESORCE): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2017 Jan 7;389(10064):56-66.
- 37e. Abou-Alfa GK, Meyer T, Cheng AL, et al. Cabozantinib in Patients with Advanced and Progressing Hepatocellular Carcinoma. *N Engl J Med*. 2018 Jul 5;379(1):54-63.
- 38e. Zhu AX, Park JO, Ryoo BY, et al. Ramucirumab versus placebo as second-line treatment in patients with advanced hepatocellular carcinoma following first-line therapy with sorafenib (REACH): a randomised, double-blind, multicentre, phase 3 trial. *Lancet Oncol*. 2015 Jul;16(7):859-70.
- 39e. Zhu AX, Kang YK, Yen CJ, et al. Ramucirumab after sorafenib in patients with advanced hepatocellular carcinoma and increased α -fetoprotein concentrations (REACH-2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol*. 2019 Feb;20(2):282-296.
- 40e. Yau T, et al. Nivolumab (NIVO) + ipilimumab (IPI) combination therapy in patients (pts) with advanced hepatocellular carcinoma (aHCC): Results from CheckMate 040 (abstract). *J Clin Oncol* 37, 2019 (suppl; abstr 4012). Abstract available inglibrary.asco.org/record/173194/abstract (Accessed on April 24, 2020).
- 41e. Paz-Ares L, et al. Pembrolizumab plus Chemotherapy for Squamous Non–Small-Cell Lung Cancer. *N Engl J Med* 2018; 379:2040-2051.
- 42e. Reck M, Rodríguez-Abreu D, Robinson AG, et al. Pembrolizumab versus Chemotherapy for PD-L1–Positive Non–Small-Cell Lung Cancer. *N Engl J Med* 2016; 375:1823-1833.
- 43e. Gandhi L, et al. Pembrolizumab plus Chemotherapy in Metastatic Non–Small-Cell Lung Cancer. *N Engl J Med* 2018; 378:2078-2092.

- 44e. Planchard D, Smit EF, Groen HJM, et al. Dabrafenib plus trametinib in patients with previously untreated BRAFV600E-mutant metastatic non-small-cell lung cancer: an open-label, phase 2 trial. *Lancet Oncol.* 2017 Oct;18(10):1307-1316. doi: 10.1016/S1470-2045(17)30679-4.
- 45e. Planchard D, Kim TM, Mazieres J, et al. Dabrafenib in patients with BRAF(V600E)-positive advanced non-small-cell lung cancer: a single-arm, multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2016 May;17(5):642-50. doi: 10.1016/S1470-2045(16)00077-2.
- 46e. Gautschi O, Milia J, Cabarro B, et al. Targeted Therapy for Patients with BRAF-Mutant Lung Cancer: Results from the European EURAF Cohort. *J Thorac Oncol.* 2015 Oct;10(10):1451-7. doi: 10.1097/JTO.0000000000000625.
- 47e. Drilon A, Laetsch TW, Kummar S, et al. Efficacy of Larotrectinib in TRK Fusion-Positive Cancers in Adults and Children. *N Engl J Med.* 2018;378(8):731–739. doi:10.1056/NEJMoa1714448.
- 48e. Doebele R, Paz-Ares L, Farago AF, et al. Entrectinib in NTRK-fusion positive (NTRK-FP) non-small cell lung cancer (NSCLC): Integrated analysis of patients enrolled in three trials (STARTRK-2, STARTRK-1 and ALKA-372-001)[abstract]. *AACR Annual Meeting.* Atlanta, GA:Abstract CT131.
- 49e. Carbone DP, Reck M, Paz-Ares L, et al. First-Line Nivolumab in Stage IV or Recurrent Non-Small-Cell Lung Cancer. *N Engl J Med.* 2017;376(25):2415–2426. doi:10.1056/NEJMoa1613493.
- 50e. West H, McCleod M, Hussein M, et al. Atezolizumab in combination with carboplatin plus nab-paclitaxel chemotherapy compared with chemotherapy alone as first-line treatment for metastatic non-squamous non-small-cell lung cancer (IMpower130): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol.* 2019;20(7):924–937. doi:10.1016/S1470-2045(19)30167-6.
- 51e. Socinski MA, Jotte RM, Cappuzzo F, et al. Atezolizumab for First-Line Treatment of Metastatic Nonsquamous NSCLC. *N Engl J Med* 2018; 378:2288-2301. DOI: 10.1056/NEJMoa1716948.
- 52e. Herbst RS, Baas P, Kim DW, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet.* 2016 Apr 9;387(10027):1540-1550. doi: 10.1016/S0140-6736(15)01281-7.
- 53e. Barlesi F, Park K, Ciardiello F, et al. Primary analysis from OAK, a randomized phase III study comparing atezolizumab with docetaxel in 2L/3L NSCLC. *Ann of Oncol* 2016 Oct;27(suppl_6):LBA44_PR.
- 54e. Mok TS, Wu Y-L, Ahn M-J, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med.* 2017;376(7):629–640. doi:10.1056/NEJMoa1612674.
- 55e. Solomon BJ, Besse B, Bauer TM, et al. Lorlatinib in patients with ALK-positive non-small-cell lung cancer: results from a global phase 2 study [published correction appears in *Lancet Oncol.*

- 2019 Jan;20(1):e10]. *Lancet Oncol*. 2018;19(12):1654–1667. doi:10.1016/S1470-2045(18)30649-1.
- 56e. Ou SH, Ahn JS, De Petris L, et al. Alectinib in Crizotinib-Refractory ALK-Rearranged Non-Small-Cell Lung Cancer: A Phase II Global Study. *J Clin Oncol*. 2016;34(7):661–668. doi:10.1200/jco.2015.63.9443.
- 57e. Huber RM, Hansen KH, Paz-Ares Rodríguez L, et al. Brigatinib in Crizotinib-Refractory ALK+ NSCLC: 2-Year Follow-up on Systemic and Intracranial Outcomes in the Phase 2 ALTA Trial. *J Thorac Oncol*. 2020;15(3):404–415. doi:10.1016/j.jtho.2019.11.004.
- 58e. Shaw AT, Kim TM, Crinò L, et al. Ceritinib versus chemotherapy in patients with ALK-rearranged non-small-cell lung cancer previously given chemotherapy and crizotinib (ASCEND-5): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol*. 2017;18(7):874–886. doi:10.1016/S1470-2045(17)30339-X.
- 59e. Rini BI, Plimack ER, Stus V, et al. Pembrolizumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2019;380(12):1116–1127. doi:10.1056/NEJMoa1816714.
- 60e. Sternberg CN, Davis ID, Mardiak J, et al. Pazopanib in locally advanced or metastatic renal cell carcinoma: results of a randomized phase III trial. *J Clin Oncol*. 2010 Feb 20;28(6):1061-8.
- 61e. Sternberg CN, Hawkins RE, Wagstaff J, et al. A randomised, double-blind phase III study of pazopanib in patients with advanced and/or metastatic renal cell carcinoma: final overall survival results and safety update. *Eur J Cancer*. 2013 Apr;49(6):1287-96.
- 62e. Motzer RJ, Hutson TE, Tomczak P, et al. Sunitinib versus interferon alfa in metastatic renal-cell carcinoma. *N Engl J Med*. 2007 Jan 11;356(2):115-24.
- 63e. Choueiri TK, Halabi S, Sanford BL, et al. Cabozantinib Versus Sunitinib As Initial Targeted Therapy for Patients With Metastatic Renal Cell Carcinoma of Poor or Intermediate Risk: The Alliance A031203 CABOSUN Trial. *J Clin Oncol*. 2016;35(6):591–597.
- 64e. Hammers HJ, Plimack ER, Infante JR, et al. Safety and Efficacy of Nivolumab in Combination With Ipilimumab in Metastatic Renal Cell Carcinoma: The CheckMate 016 Study. *J Clin Oncol*. 2017;35(34):3851–3858. doi:10.1200/JCO.2016.72.1985.
- 65e. Choueiri TK, Escudier B, Powles T, et al. Cabozantinib versus Everolimus in Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2015;373(19):1814–1823. doi:10.1056/NEJMoa1510016.
- 66e. Koshkin VS, Barata PC, Zhang T, et al. Clinical activity of nivolumab in patients with non-clear cell renal cell carcinoma. *J Immunother Cancer*. 2018;6(1):9. Published 2018 Jan 29. doi:10.1186/s40425-018-0319-9.
- 67e. Armstrong AJ, Halabi S, Eisen T, et al. Everolimus versus sunitinib for patients with metastatic non-clear cell renal cell carcinoma (ASPEN): a multicentre, open-label, randomised phase 2 trial. *Lancet Oncol*. 2016;17(3):378–388. doi:10.1016/S1470-2045(15)00515-X.

- 68e. Campbell MT, Bilen MA, Shah AY, et al. Cabozantinib for the treatment of patients with metastatic non-clear cell renal cell carcinoma: A retrospective analysis. *Eur J Cancer*. 2018;104:188–194. doi:10.1016/j.ejca.2018.08.014.
- 69e. Blank CU, Bono P, Larkin JMG, et al. Safety and efficacy of everolimus in patients with non-clear cell renal cell carcinoma refractory to VEGF-targeted therapy: Subgroup analysis of REACT [abstract]. *J Clin Oncol* 2012; 30 (5_suppl):Abstract 402.
- 70e. Chen R, Zinzani PL, Fanale MA, et al. Phase II Study of the Efficacy and Safety of Pembrolizumab for Relapsed/Refractory Classic Hodgkin Lymphoma. *J Clin Oncol*. 2017;35(19):2125–2132. doi:10.1200/JCO.2016.72.1316.
- 71e. Chen R, Zinzani PL, Lee HJ, et al. Pembrolizumab in relapsed or refractory Hodgkin lymphoma: 2-year follow-up of KEYNOTE-087. *Blood*. 2019;134(14):1144–1153. doi:10.1182/blood.2019000324.
- 72e. Armand P, Shipp MA, Ribrag V, et al. Programmed Death-1 Blockade With Pembrolizumab in Patients With Classical Hodgkin Lymphoma After Brentuximab Vedotin Failure. *J Clin Oncol*. 2016;34(31):3733–3739. doi:10.1200/JCO.2016.67.3467.
- 73e. Younes A, Santoro A, Shipp M, et al. Nivolumab for classical Hodgkin's lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: a multicentre, multicohort, single-arm phase 2 trial. *Lancet Oncol*. 2016;17(9):1283–1294. doi:10.1016/S1470-2045(16)30167-X.
- 74e. Moskowitz AJ, Hamlin PA Jr, Perales MA, et al. Phase II study of bendamustine in relapsed and refractory Hodgkin lymphoma. *J Clin Oncol*. 2013;31(4):456–460. doi:10.1200/JCO.2012.45.3308.
- 75e. Gopal AK, Press OW, Shustov AR, et al. Efficacy and safety of gemcitabine, carboplatin, dexamethasone, and rituximab in patients with relapsed/refractory lymphoma: a prospective multi-center phase II study by the Puget Sound Oncology Consortium. *Leuk Lymphoma*. 2010;51(8):1523–1529. doi:10.3109/10428194.2010.491137.
- 76e. Rodriguez MA, Cabanillas FC, Hagemeister FB, et al. A phase II trial of mesna/ifosfamide, mitoxantrone and etoposide for refractory lymphomas. *Ann Oncol*. 1995 Jul;6(6):609-11.
- 77e. Colwill R, Crump M, Couture F, et al. Mini-BEAM as salvage therapy for relapsed or refractory Hodgkin's disease before intensive therapy and autologous bone marrow transplantation. *J Clin Oncol*. 1995 Feb;13(2):396-402.
- 78e. Martín A, Fernández-Jiménez MC, et al. Long-term follow-up in patients treated with Mini-BEAM as salvage therapy for relapsed or refractory Hodgkin's disease. *Br J Haematol*. 2001 Apr;113(1):161-71.
- 79e. Herrera AF, Moskowitz AJ, Bartlett NL, et al. Interim results of brentuximab vedotin in combination with nivolumab in patients with relapsed or refractory Hodgkin lymphoma. *Blood*. 2018;131(11):1183–1194. doi:10.1182/blood-2017-10-811224.

- 80e. Chen RW, Palmer J, Martin, et al. Results of a Phase II Trial of Brentuximab Vedotin As First Line Salvage Therapy in Relapsed/Refractory HL Prior to AHCT [abstract]. *Blood* 2014;124:Abstract 501.
- 81e. O'Connor OA, Lue JK, Sawas A, et al. Brentuximab vedotin plus bendamustine in relapsed or refractory Hodgkin's lymphoma: an international, multicentre, single-arm, phase 1-2 trial. *Lancet Oncol* 2018; 19:257.
- 82e. Cohen EEW, Soulières D, Le Tourneau C, et al. Pembrolizumab versus methotrexate, docetaxel, or cetuximab for recurrent or metastatic head-and-neck squamous cell carcinoma (KEYNOTE-040): a randomised, open-label, phase 3 study. *Lancet*. 2019 Jan 12;393(10167):156-167.
- 83e. Cohen EE, Harrington KJ, Le Tourneau C, et al. LBA45_PR - Pembrolizumab (pembro) vs standard of care (SOC) for recurrent or metastatic head and neck squamous cell carcinoma (R/M HNSCC): Phase 3 KEYNOTE-040. *Ann Oncol*. 2017;28(suppl_5):605-649. doi:10.1093/annonc/mdx440.
- 84e. Chow LQM, Haddad R, Gupta S, et al. Antitumor Activity of Pembrolizumab in Biomarker-Unselected Patients With Recurrent and/or Metastatic Head and Neck Squamous Cell Carcinoma: Results From the Phase Ib KEYNOTE-012 Expansion Cohort. *J Clin Oncol*. 2016;34(32):3838–3845. doi:10.1200/JCO.2016.68.1478.
- 85e. Bellmunt J, de Wit R, Vaughn DJ, et al. Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma. *N Engl J Med*. 2017;376(11):1015–1026. doi:10.1056/NEJMoa1613683.
- 86e. Apolo AB, Infante JR, Balmanoukian A, et al. Avelumab, an Anti-Programmed Death-Ligand 1 Antibody, In Patients With Refractory Metastatic Urothelial Carcinoma: Results From a Multicenter, Phase Ib Study. *J Clin Oncol*. 2017;35(19):2117–2124. doi:10.1200/JCO.2016.71.6795.
- 87e. Patel MR, Ellerton J, Infante JR, et al. Avelumab in metastatic urothelial carcinoma after platinum failure (JAVELIN Solid Tumor): pooled results from two expansion cohorts of an open-label, phase 1 trial. *Lancet Oncol*. 2018 Jan;19(1):51-64.
- 88e. Rosenberg JE, Hoffman-Censits J, Powles T, et al. Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet*. 2016;387(10031):1909–1920. doi:10.1016/S0140-6736(16)00561-4.
- 89e. Powles T, Durán I, van der Heijden MS, et al. Atezolizumab versus chemotherapy in patients with platinum-treated locally advanced or metastatic urothelial carcinoma (IMvigor211): a multicentre, open-label, phase 3 randomised controlled trial. *Lancet*. 2018 Feb 24;391(10122):748-757. doi: 10.1016/S0140-6736(17)33297-X.
- 90e. Massard C, Gordon MS, Sharma S, et al. Safety and Efficacy of Durvalumab (MEDI4736), an Anti-Programmed Cell Death Ligand-1 Immune Checkpoint Inhibitor, in Patients With Advanced

Urothelial Bladder Cancer. *J Clin Oncol*. 2016;34(26):3119–3125.
doi:10.1200/JCO.2016.67.9761.

- 91e. Siefker-Radtke AO, Necchi A, Park SH, et al. First results from the primary analysis population of the phase 2 study of erdafitinib (ERDA; JNJ-42756493) in patients (pts) with metastatic or unresectable urothelial carcinoma (mUC) and FGFR alterations (FGFRalt). *J Clin Oncol* 2018;36(15_suppl):4503.
- 92e. Chung HC, Piha-Paul SA, Lopez-Martin J, et al. Pembrolizumab After Two or More Lines of Previous Therapy in Patients With Recurrent or Metastatic SCLC: Results From the KEYNOTE-028 and KEYNOTE-158 Studies. *J Thorac Oncol*. 2020;15(4):618–627.
doi:10.1016/j.jtho.2019.12.109.
- 93e. Chung HC, Lopez-Martin JA, Kao S C-H, et al. Phase 2 study of pembrolizumab in advanced small-cell lung cancer (SCLC): KEYNOTE-158. *J Clin Oncol* 2018; 36S: ASCO# 8506.
- 94e. Ott PA, Elez E, Hirt S, et al. Pembrolizumab in Patients With Extensive-Stage Small-Cell Lung Cancer: Results From the Phase Ib KEYNOTE-028 Study. *J Clin Oncol* 2017; 35:3823.
- 95e. Reck M, Vicente D, Ciuleanu T, et al. LBA5: Efficacy and safety of nivolumab (nivo) monotherapy versus chemotherapy (chemo) in recurrent small cell lung cancer (SCLC): Results from CheckMate 331 [abstract]. *Ann Oncol* 2018;29:43.
- 96e. von Pawel J, Schiller JH, Shepherd FA, et al. Topotecan versus cyclophosphamide, doxorubicin, and vincristine for the treatment of recurrent small-cell lung cancer. *J Clin Oncol*. 1999 Feb;17(2):658-67.
- 97e. O'Brien ME, Ciuleanu TE, Tsekov H, et al. Phase III trial comparing supportive care alone with supportive care with oral topotecan in patients with relapsed small-cell lung cancer. *J Clin Oncol*. 2006 Dec 1;24(34):5441-7.
- 98e. Yamamoto N, Tsurutani J, Yoshimura N, et al. Phase II study of weekly paclitaxel for relapsed and refractory small cell lung cancer. *Anticancer Res*. 2006 Jan-Feb;26(1B):777-81.
- 99e. Smit EF, Fokkema E, Biesma B, Groen HJ, Snoek W, Postmus PE. A phase II study of paclitaxel in heavily pretreated patients with small-cell lung cancer. *Br J Cancer*. 1998;77(2):347–351. doi:10.1038/bjc.1998.54.
- 100e. Smyth JF, Smith IE, Sessa C, et al. Activity of docetaxel (Taxotere) in small cell lung cancer. The Early Clinical Trials Group of the EORTC. *Eur J Cancer*. 1994;30A(8):1058-60.
- 101e. Masuda N, Fukuoka M, Kusunoki Y, et al. CPT-11: a new derivative of camptothecin for the treatment of refractory or relapsed small-cell lung cancer. *J Clin Oncol*. 1992 Aug;10(8):1225-9.
- 102e. Pietanza MC, Kadota K, Huberman K, et al. Phase II trial of temozolomide in patients with relapsed sensitive or refractory small cell lung cancer, with assessment of methylguanine-DNA methyltransferase as a potential biomarker. *Clin Cancer Res*. 2012 Feb 15;18(4):1138-45. doi: 10.1158/1078-0432.CCR-11-2059.

- 103e. Lenz HJ, Lonardi S, Zagonel V, et al. Nivolumab (NIVO) + low-dose ipilimumab (IPI) as first-line (1L) therapy in microsatellite instability-high/DNA mismatch repair deficient (MSI-H/dMMR) metastatic colorectal cancer (mCRC): Clinical update [abstract]. *Journal of Clinical Oncology* 2019;37:3521-3521.
- 104e. Le DT, Uram JN, Wang H, et al. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med*. 2015;372(26):2509–2520. doi:10.1056/NEJMoa1500596.
- 105e. Le DT, Kim TW, Van Cutsem E, et al. Phase II Open-Label Study of Pembrolizumab in Treatment-Refractory, Microsatellite Instability-High/Mismatch Repair-Deficient Metastatic Colorectal Cancer: KEYNOTE-164. *J Clin Oncol* 2020; 38:11.
- 106e. Nghiem PT, Bhatia S, Lipson EJ, et al. PD-1 Blockade with Pembrolizumab in Advanced Merkel-Cell Carcinoma. *N Engl J Med*. 2016;374(26):2542–2552. doi:10.1056/NEJMoa1603702.
- 107e. D'Angelo SP, Russell J, Lebbé C, et al. Efficacy and Safety of First-line Avelumab Treatment in Patients With Stage IV Metastatic Merkel Cell Carcinoma: A Preplanned Interim Analysis of a Clinical Trial. *JAMA Oncol*. 2018;4(9):e180077. doi:10.1001/jamaoncol.2018.0077.
- 108e. Kaufman HL, Russell JS, Hamid O, et al. Updated efficacy of avelumab in patients with previously treated metastatic Merkel cell carcinoma after ≥1 year of follow-up: JAVELIN Merkel 200, a phase 2 clinical trial. *J Immunother Cancer*. 2018;6(1):7. Published 2018 Jan 19. doi:10.1186/s40425-017-0310-x.
- 109e. Kluger HM, Chiang V, Mahajan A, et al. Long-Term Survival of Patients With Melanoma With Active Brain Metastases Treated With Pembrolizumab on a Phase II Trial. *J Clin Oncol*. 2019;37(1):52–60. doi:10.1200/JCO.18.00204.
- 110e. Margolin K, Ernstoff MS, Hamid O, et al. Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial. *Lancet Oncol*. 2012 May;13(5):459-65. doi: 10.1016/S1470-2045(12)70090-6.
- 111e. Goldberg SB, Gettinger SN, Mahajan A, et al. Pembrolizumab for patients with melanoma or non-small-cell lung cancer and untreated brain metastases: early analysis of a non-randomised, open-label, phase 2 trial. *Lancet Oncol*. 2016;17(7):976–983. doi:10.1016/S1470-2045(16)30053-5.
- 112e. Ott PA, Piha-Paul SA, Munster P, et al. Safety and antitumor activity of the anti-PD-1 antibody pembrolizumab in patients with recurrent carcinoma of the anal canal. *Ann Oncol*. 2017;28(5):1036–1041. doi:10.1093/annonc/mdx029.
- 113e. Ghorani E, Kaur B, Fisher RA, et al. Pembrolizumab is effective for drug-resistant gestational trophoblastic neoplasia. *Lancet* 2017; 390:2343.
- 114e. Alley EW, Lopez J, Santoro A, et al. Clinical safety and activity of pembrolizumab in patients with malignant pleural mesothelioma (KEYNOTE-028): preliminary results from a non-randomised, open-label, phase 1b trial. *Lancet Oncol*. 2017 May;18(5):623-630.

- 115e. Alley EW, Lopez J, Santoro A, et al. Long-Term Overall Survival for Patients with Malignant Pleural Mesothelioma on Pembrolizumab Enrolled in KEYNOTE-028. *J Thorac Oncol*. 2017 Jan;12(1):S294.
- 116e. Metaxas Y, Rivalland G, Mauti LA, et al. Pembrolizumab as Palliative Immunotherapy in Malignant Pleural Mesothelioma. *J Thorac Oncol*. 2018 Nov;13(11):1784-1791.
- 117e. Jassem J, Ramlau R, Santoro A, et al, "Phase III Trial of Pemetrexed Plus Best Supportive Care Compared With Best Supportive Care in Previously Treated Patients With Advanced Malignant Pleural Mesothelioma," *J Clin Oncol*, 2008, 26(10):1698-704. [PubMed 18375898]
- 118e. Zucali PA, Simonelli M, Michetti G, et al. Second-line chemotherapy in malignant pleural mesothelioma: results of a retrospective multicenter survey. *Lung Cancer*. 2012 Mar;75(3):360-7.
- 119e. Quispel-Janssen J, van der Noort V, de Vries JF, et al. Programmed Death 1 Blockade With Nivolumab in Patients With Recurrent Malignant Pleural Mesothelioma. *J Thorac Oncol*. 2018 Oct;13(10):1569-1576.
- 120e. Popat S, Curioni-Fontecedro A, Polydoropoulou V, et al. A multicentre randomized phase III trial comparing pembrolizumab (P) versus single-agent chemotherapy (CT) for advanced pretreated malignant pleural mesothelioma (MPM): Results from the European Thoracic Oncology Platform (ETOP 9-15) PROMISE-meso trial. *Ann Oncol* 2019; 30S: ESMO #LBA91_PR.
- 121e. Chung HC, Ros W, Derlord JP, et al. Efficacy and Safety of Pembrolizumab in Previously Treated Advanced Cervical Cancer: Results From the Phase II KEYNOTE-158 Study. *J Clin Oncol*. 2019;37(17):1470-1478.
- 122e. Kwong YL, Chan TSY, Tan D, et al. PD1 blockade with pembrolizumab is highly effective in relapsed or refractory NK/T-cell lymphoma failing l-asparaginase. *Blood* 2017; 129:2437.
- 123e. Rittmeyer A, Barlesi F, Waterkamp D, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial [published correction appears in *Lancet*. 2017 Apr 8;389(10077):e5]. *Lancet*. 2017;389(10066):255-265. doi:10.1016/S0140-6736(16)32517-X.
- 124e. Gauvain C, Vauléon E, Chouaid C, et al. Intracerebral efficacy and tolerance of nivolumab in non-small-cell lung cancer patients with brain metastases [published correction appears in *Lung Cancer*. 2019 Oct;136:159]. *Lung Cancer*. 2018;116:62-66. doi:10.1016/j.lungcan.2017.12.008.
- 125e. Rizvi NA, Mazières J, Planchard D, et al. Activity and safety of nivolumab, an anti-PD-1 immune checkpoint inhibitor, for patients with advanced, refractory squamous non-small-cell lung cancer (CheckMate 063): a phase 2, single-arm trial. *Lancet Oncol*. 2015;16(3):257-265. doi:10.1016/S1470-2045(15)70054-9.

- 126e. Herbst RS, Giaccone G, de Marinis F, et al. Atezolizumab for First-Line Treatment of PD-L1-Selected Patients with NSCLC. *N Engl J Med*. 2020 Oct 1;383(14):1328-1339. doi: 10.1056/NEJMoa1917346.
- 127e. Assersohn L, Brown G, Cunningham D, et al. Phase II study of irinotecan and 5-fluorouracil/leucovorin in patients with primary refractory or relapsed advanced oesophageal and gastric carcinoma. *Ann Oncol*. 2004;15(1):64-69. doi:10.1093/annonc/mdh007.
- 128e. Cole PD, Mauz-Körholz C, Mascarini, M, et al. Nivolumab and brentuximab vedotin (BV)-based, response-adapted treatment in children, adolescents, and young adults (CAYA) with standard-risk relapsed/refractory classical Hodgkin lymphoma (R/R cHL): Primary analysis. *Journal of Clinical Oncology* 2020;38:8013.
- 129e. Cole PD, McCarten KM, Pei Q, et al. Brentuximab vedotin with gemcitabine for paediatric and young adult patients with relapsed or refractory Hodgkin's lymphoma (AHOD1221): a Children's Oncology Group, multicentre single-arm, phase 1-2 trial. *Lancet Oncol*. 2018 Sep;19(9):1229-1238. doi: 10.1016/S1470-2045(18)30426-1.
- 130e. Kelly R, Ajani J, Kuzdzal J, et al. Adjuvant nivolumab in resected esophageal or gastroesophageal junction cancer (EC/GEJC) following neoadjuvant chemoradiation therapy (CRT): First results of the CheckMate 577 study. [abstract]. Presented at the Oral Presentation presented at the ESMO 2020 Annual Meeting; September 19-21, 2020; Virtual Meeting.
- 131e. Kato K, Sun JM, Shah M.A., et al. Pembrolizumab plus chemotherapy versus chemotherapy as first-line therapy in patients with advanced esophageal cancer: The phase 3 KEYNOTE-590 study. [abstract]. Presented at the Oral Presentation presented at the ESMO 2020 Annual Meeting; September 19-21, 2020; Virtual Meeting.
- 132e. Kojima T, Shah MA, Muro K, et al. Randomized Phase III KEYNOTE-181 Study of Pembrolizumab Versus Chemotherapy in Advanced Esophageal Cancer. *J Clin Oncol*. 2020 Dec 10;38(35):4138-4148. doi: 10.1200/JCO.20.01888.
- 133e. Andre T, Shiu KK, Kim TW, et al. Pembrolizumab versus chemotherapy for microsatellite instability-high/mismatch repair deficient metastatic colorectal cancer: The phase 3 KEYNOTE-177 Study. *J Clin Oncol*. 2020;38(18_suppl):LBA4-LBA4.
- 134e. Goldman JW, Crino L, Vokes EE, et al. Nivolumab (nivo) in Patients (pts) With Advanced (adv) NSCLC and Central Nervous System (CNS) Metastases (mets). *J Thorac Oncol* 2016;11:S238-S239.
- 135e. Zalcman G, Mazieres J, Margery J, et al. Bevacizumab for newly diagnosed pleural mesothelioma in the Mesothelioma Avastin Cisplatin Pemetrexed Study (MAPS): a randomised, controlled, open-label, phase 3 trial. *Lancet*. 2016 Apr 2;387(10026):1405-1414. doi: 10.1016/S0140-6736(15)01238-6. Epub 2015 Dec 21.

- 136e. Vogelzang NJ, Rusthoven JJ, Symanowski J, et al. Phase III study of pemetrexed in combination with cisplatin versus cisplatin alone in patients with malignant pleural mesothelioma. *J Clin Oncol*. 2003 Jul 15;21(14):2636-44. doi: 10.1200/JCO.2003.11.136.
- 137e. Motzer R, Alekseev B, Rha SY, et al. Lenvatinib plus Pembrolizumab or Everolimus for Advanced Renal Cell Carcinoma. *N Engl J Med*. 2021 Apr 8;384(14):1289-1300. doi: 10.1056/NEJMoa2035716.
- 138e. Yau T, Park JW, Finn RS, et al. LBA38_PR - CheckMate 459: A randomized, multi-center phase III study of nivolumab (NIVO) vs sorafenib (SOR) as first-line (1L) treatment in patients (pts) with advanced hepatocellular carcinoma (aHCC). *Ann of Oncol* 2019 Oct;30(suppl_5):v874-v875.
- 139e. Powles T, Rosenberg JE, Sonpavde GP, et al. Enfortumab Vedotin in Previously Treated Advanced Urothelial Carcinoma. *N Engl J Med*. 2021 Mar 25;384(12):1125-1135. doi: 10.1056/NEJMoa2035807. Epub 2021 Feb 12.
- 140e. Balar AV, McGregor BA, Rosenberg JE, et al. EV-201 Cohort 2: Enfortumab vedotin in cisplatin-ineligible patients with locally advanced or metastatic urothelial cancer who received prior PD-1/PD-L1 inhibitors. DOI: 10.1200/JCO.2021.39.6_suppl.394 *Journal of Clinical Oncology* 39, no. 6_suppl (February 20, 2021) 394-394.
- 141e. Sternberg CN, de Mulder PH, Schornagel JH, et al. Randomized phase III trial of high-dose-intensity methotrexate, vinblastine, doxorubicin, and cisplatin (MVAC) chemotherapy and recombinant human granulocyte colony-stimulating factor versus classic MVAC in advanced urothelial tract tumors: European Organization for Research and Treatment of Cancer Protocol no. 30924. *J Clin Oncol*. 2001 May 15;19(10):2638-46. doi: 10.1200/JCO.2001.19.10.2638.
- 142e. von der Maase H, Hansen SW, Roberts JT, et al. Gemcitabine and cisplatin versus methotrexate, vinblastine, doxorubicin, and cisplatin in advanced or metastatic bladder cancer: results of a large, randomized, multinational, multicenter, phase III study. *J Clin Oncol*. 2000 Sep;18(17):3068-77. doi: 10.1200/JCO.2000.18.17.3068.
- 143e. Zalcman G, Mazieres J, Greillier L, et al. Second/third-line nivolumab vs nivo plus ipilimumab in malignant pleural mesothelioma: Long-term results of IFCT-1501 MAPS2 phase IIR trial with a focus on hyperprogression (HPD). *Ann of Oncol* 2016 Oct;30(suppl_5):v747.
- 144e. Sosman JA. (2021). Immunotherapy of advanced melanoma with immune checkpoint inhibition. In Atkins MB & Shah S (Eds.), UpToDate. Availthttps://www.uptodate.com/contents/immunotherapy-of-advanced-melanoma-with-immune-checkpoint-inhibition?search=melanoma%20treatment&topicRef=85841&source=see_link#H177442841.
- 145e. Motzer RJ, Hutson TE, Glen H, et al. Lenvatinib, everolimus, and the combination in patients with metastatic renal cell carcinoma: a randomised, phase 2, open-label, multicentre trial. *Lancet Oncol*. 2015 Nov;16(15):1473-1482. doi: 10.1016/S1470-2045(15)00290-9. Epub 2015

Oct 22. Erratum in: *Lancet Oncol*. 2016 Jul;17 (7):e270. Erratum in: *Lancet Oncol*. 2018 Oct;19(10):e509.

- 146e. Sezer A, Kilickap S, Gümüş M, et al. Cemiplimab monotherapy for first-line treatment of advanced non-small-cell lung cancer with PD-L1 of at least 50%: a multicentre, open-label, global, phase 3, randomised, controlled trial. *Lancet*. 2021 Feb 13;397(10274):592-604. doi: 10.1016/S0140-6736(21)00228-2.
- 147e. Tawbi HA, Forsyth PA, Algazi A, et al. Combined Nivolumab and Ipilimumab in Melanoma Metastatic to the Brain. *N Engl J Med*. 2018;379(8):722-730. doi:10.1056/NEJMoa1805453.
- 148e. Spigel DR, Vicente D, Ciuleanu TE, et al. Second-line nivolumab in relapsed small-cell lung cancer: CheckMate 331☆. *Ann Oncol*. 2021 May;32(5):631-641. doi: 10.1016/j.annonc.2021.01.071. Epub 2021 Feb 1.
- 149e. Wolchok JD, Chiarion-Sileni V, Gonzalez R, et al. CheckMate 067: 6.5-year outcomes in patients (pts) with advanced melanoma. *J Clin Oncol* 2021; 39;15S.
- 150e. Pires da Silva I, Ahmed T, Reijers ILM, et al. Ipilimumab alone or ipilimumab plus anti-PD-1 therapy in patients with metastatic melanoma resistant to anti-PD-(L)1 monotherapy: a multicentre, retrospective, cohort study. *Lancet Oncol*. 2021 Jun;22(6):836-847. doi: 10.1016/S1470-2045(21)00097-8.
- 151e. Chung HC, Ros W, Delord JP, et al. Efficacy and Safety of Pembrolizumab in Previously Treated Advanced Cervical Cancer: Results From the Phase II KEYNOTE-158 Study. *J Clin Oncol*. 2019 Jun 10;37(17):1470-1478. doi: 10.1200/JCO.18.01265. Epub 2019 Apr 3.
- 152e. Olson D, Luke J, Poklepovic A, et al. Significant antitumor activity for low-dose ipilimumab (IPI) with pembrolizumab (PEMBRO) immediately following progression on PD1 Ab in melanoma (MEL) in a phase II trial. *Journal of Clinical Oncology* 2020 38:15_suppl, 10004-10004.
- 153e. Berton D, Banerjee S, Curigliano G, et al. Antitumor activity of dostarlimab in patients with mismatch repair-deficient/microsatellite instability–high tumors: A combined analysis of two cohorts in the GARNET study. *Journal of Clinical Oncology*. Volume 39, Issue 15_suppl. doi/abs/10.1200/JCO.2021.39.15_suppl.2564.
- 154e. Baas P, Scherpereel A, Nowak AK, et al. First-line nivolumab plus ipilimumab in unresectable malignant pleural mesothelioma (CheckMate 743): a multicentre, randomised, open-label, phase 3 trial. *Lancet*. 2021 Jan 30;397(10272):375-386. doi: 10.1016/S0140-6736(20)32714-8. Epub 2021 Jan 21. Erratum in: *Lancet*. 2021 Feb 20;397(10275):670.
- 155e. Kato K, Shah MA, Enzinger P, et al. KEYNOTE-590: Phase III study of first-line chemotherapy with or without pembrolizumab for advanced esophageal cancer. *Future Oncol*. 2019 Apr;15(10):1057-1066. doi: 10.2217/fon-2018-0609.

- 156e. Motzer RJ, Penkov K, Haanen J, et al. Avelumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2019 Mar 21;380(12):1103-1115. doi: 10.1056/NEJMoa1816047. Epub 2019 Feb 16.
- 157e. Tawbi HA, Schadendorf D, Lipson EJ, et al. Relatlimab and Nivolumab versus Nivolumab in Untreated Advanced Melanoma. *N Engl J Med*. 2022 Jan 6;386(1):24-34. doi: 10.1056/NEJMoa2109970.
- 158e. Vanderwalde AM, Moon J, Kendra K et al. S1616: Ipilimumab plus nivolumab versus ipilimumab alone in patients with metastatic or unresectable melanoma that did not respond to anti-PD-1 therapy In: Proceedings of the 113th Annual Meeting of the American Association for Cancer Research; 2021 April 8-13; New Orleans LA. Philadelphia (PA): AACR; 2022. Abstract CT013. <https://www.abstractsonline.com/pp8/#!/10517/presentation/20155> (Accessed on June 10, 2022).
- 159e. Forde PM, Spicer J, Lu S, et al. Neoadjuvant Nivolumab plus Chemotherapy in Resectable Lung Cancer. *N Engl J Med*. 2022 May 26;386(21):1973-1985. doi: 10.1056/NEJMoa2202170. Epub 2022 Apr 11.
- 160e. Raghav KPS, Overman MJ, Liu S, et al. A phase II trial of atezolizumab and bevacizumab in patients with relapsed/refractory and unresectable malignant peritoneal mesothelioma. *Journal of Clinical Oncology* 2020 38:15_suppl, 9013-9013.
- 161e. Lee CH, Voss MH, Carlo MI, et al. Phase II Trial of Cabozantinib Plus Nivolumab in Patients With Non-Clear-Cell Renal Cell Carcinoma and Genomic Correlates. *J Clin Oncol*. 2022 Jul 20;40(21):2333-2341.
- 162e. Atkins MB, Lee SJ, Chmielowski B, et al. DREAMseq (Doublet, Randomized Evaluation in Advanced Melanoma Sequencing): A phase III trial—ECOG-ACRIN EA6134. *J Clin Oncol*. 2021 Dec 20;39(36_suppl):356154-356-154.
- 163e. Mok TS, Wu Y-L, Ahn M-J, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med*. 2017 Feb 16;376(7):629-640.
- 164e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Pediatric Central Nervous System Cancers. Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 165e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Uterine Neoplasms. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are

trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.

- 166e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Vulvar Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2026.
- 167e. André T, Lonardi S, Wong KYM, et al. Nivolumab plus low-dose ipilimumab in previously treated patients with microsatellite instability-high/mismatch repair-deficient metastatic colorectal cancer: 4-year follow-up from CheckMate 142. *Ann Oncol*. 2022 Oct;33(10):1052-1060.
- 168e. Johnson ML, Cho BC, Luft A, et al; POSEIDON investigators. Durvalumab With or Without Tremelimumab in Combination With Chemotherapy as First-Line Therapy for Metastatic Non-Small-Cell Lung Cancer: The Phase III POSEIDON Study. *J Clin Oncol*. 2022 Nov 3;JCo2200975. doi: 10.1200/JCO.22.00975.
- 169e. Gogishvili M, Melkadze T, Makharadze T, et al. LBA51 EMPOWER-Lung 3: Cemiplimab in combination with platinum doublet chemotherapy for first-line (1L) treatment of advanced non-small cell lung cancer (NSCLC). *Annals of Oncology*, ISSN: 0923-7534, Vol: 32, SUPPLEMENT 5, S1328, SEPTEMBER 01, 2021. DOI10.1016/j.annonc.2021.08.2130.
- 170e. Delyon J, Biard L, Renaud M, et al. PD-1 blockade with pembrolizumab in classic or endemic Kaposi's sarcoma: a multicentre, single-arm, phase 2 study. *Lancet Oncol*. 2022 Apr;23(4):491-500. doi: 10.1016/S1470-2045(22)00097-3.
- 171e. Zer A, Licht O, Yosef L, et al. Phase II single-arm study of nivolumab and ipilimumab (Nivo/Ipi) in previously treated classical Kaposi sarcoma (cKS). *Ann Oncol*. 2022 Jul;33(7):720-727. doi: 10.1016/j.annonc.2022.03.012.
- 172e. Rao S, Guren MG, Khan K, et al. Anal cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2021 Sep;32(9):1087-1100. doi: 10.1016/j.annonc.2021.06.015.
- 173e. Marabelle A, Fakih M, Lopez J, et al. Association of Tumor Mutational Burden with Outcomes in Patients with Select Advanced Solid Tumors Treated with Pembrolizumab in KEYNOTE-158. *Ann Oncol*. 2019;30(suppl_5):v475-v532. doi: 10.1093/annonc/mdz253.
- 174e. Polizzotto MN, Uldrick TS, Wyvill KM, et al. Pomalidomide for Symptomatic Kaposi's Sarcoma in People With and Without HIV Infection: A Phase I/II Study. *J Clin Oncol*. 2016 Dec;34(34):4125-4131.
- 175e. Marabelle A, Cassier PA, Fakih M, et al. Pembrolizumab for previously treated advanced anal squamous cell carcinoma: results from the non-randomised, multicohort, multicentre, phase 2 KEYNOTE-158 study. *Lancet Gastroenterol Hepatol*. 2022 May;7(5):446-454.

- 176e. Ludford K, Ho WJ, Thomas JV, et al. Neoadjuvant Pembrolizumab in Localized Microsatellite Instability High/Deficient Mismatch Repair Solid Tumors. *J Clin Oncol*. 2023 Apr 20;41(12):2181-2190.
- 177e. Pietrantonio F, Raimondi A, Lonardi S, et al. INFINITY: A multicentre, single-arm, multi-cohort, phase II trial of tremelimumab and durvalumab as neoadjuvant treatment of patients with microsatellite instability-high (MSI) resectable gastric or gastroesophageal junction adenocarcinoma (GAC/GEJAC). *Journal of Clinical Oncology* 2023;41:358-358.
- 178e. Andre T, Elez E, Van Cutsem E, Jensen LH. Nivolumab (NIVO) plus ipilimumab (IPI) vs chemotherapy (chemo) as first-line (1L) treatment for microsatellite instability-high/mismatch repair-deficient (MSI-H/dMMR) metastatic colorectal cancer (mCRC): First results of the CheckMate 8HW study. *J Clin Oncol* 2024; 42:3S.
- 179e. Qin S, Chen Z, Fang W, et al. Pembrolizumab Versus Placebo as Second-Line Therapy in Patients From Asia With Advanced Hepatocellular Carcinoma: A Randomized, Double-Blind, Phase III Trial. *J Clin Oncol*. 2023 Mar 1;41(7):1434-1443.
- 180e. Rozeman EA, Menzies AM, van Akkooi ACJ, et al. Identification of the optimal combination dosing schedule of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma (OpACIN-neo): a multicentre, phase 2, randomised, controlled trial. *Lancet Oncol*. 2019 Jul;20(7):948-960.
- 181e. Patel SP, Othus M, Chen Y, et al. Neoadjuvant-Adjuvant or Adjuvant-Only Pembrolizumab in Advanced Melanoma. *N Engl J Med*. 2023 Mar 2;388(9):813-823.
- 182e. Zhou C, Tang KJ, Cho BC, et al. Amivantamab plus Chemotherapy in NSCLC with EGFR Exon 20 Insertions. *N Engl J Med*. 2023 Nov 30;389(22):2039-2051.
- 183e. Powles TB, Valderrama B, Gupta S, et al. LBA6 EV-302/KEYNOTE-A39: Open-label, randomized phase III study of enfortumab vedotin in combination with pembrolizumab (EV +P) vs chemotherapy in previously untreated locally advanced metastatic urothelial carcinoma (la/mUC). *Annals of Oncology*, Volume 34, Issue Supplement_2, October 2023, MDZ250.002, <https://doi.org/10.1016/j.annonc.2023.10.106>.
- 184e. De Santis M, Bellmunt J, Mead G, et al. Randomized phase II/III trial assessing gemcitabine/carboplatin and methotrexate/carboplatin/vinblastine in patients with advanced urothelial cancer who are unfit for cisplatin-based chemotherapy: EORTC study 30986. *J Clin Oncol*. 2012;30(2):191–199. doi:10.1200/JCO.2011.37.3571.
- 185e. Powles T, Park SH, Voog E, et al. Maintenance avelumab + best supportive care (BSC) versus BSC alone after platinum-based first-line (1L) chemotherapy in advanced urothelial carcinoma (UC): JAVELIN Bladder 100 phase III interim analysis. *J Clin Oncol*. 2020;38(18_suppl):LBA1-LBA1.

- 186e. Novello S, Mazières J, Oh IJ, et al. Alectinib versus chemotherapy in crizotinib-pretreated anaplastic lymphoma kinase (ALK)-positive non-small-cell lung cancer: results from the phase III ALUR study. *Ann Oncol.* 2018;29(6):1409-1416. doi:10.1093/annonc/mdy121.
- 187e. Kim DW, Tiseo M, Ahn MJ, et al. Brigatinib in Patients With Crizotinib-Refractory Anaplastic Lymphoma Kinase-Positive Non-Small-Cell Lung Cancer: A Randomized, Multicenter Phase II Trial. *J Clin Oncol.* 2017 Aug 1;35(22):2490-2498. doi: 10.1200/JCO.2016.71.5904.
- 188e. Shaw AT, Kim DW, Nakagawa K, et al. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med.* 2013 Jun 20;368(25):2385-94. doi: 10.1056/NEJMoa1214886.
- 189e. Grignani G, Rutkowski P, Lebbé C. 1146P Updated results from POD1UM-201: A phase II study of retifanlimab in patients with advanced or metastatic Merkel cell carcinoma (MCC). Presented at the European Society for Medical Oncology Congress 2023. Madrid, Spain; October 20-24, 2023
- 190e. Marabelle A, Le DT, Ascierto PA, et al. Efficacy of Pembrolizumab in Patients With Noncolorectal High Microsatellite Instability/Mismatch Repair–Deficient Cancer: Results From the Phase II KEYNOTE-158 Study. *Journal of Clinical Oncology.* 2020;38(1):1-10. doi:https://doi.org/10.1200/jco.19.02105
- 191e. Oaknin A, Tinker AV, Gilbert L, et al. Clinical Activity and Safety of the Anti–Programmed Death 1 Monoclonal Antibody Dostarlimab for Patients With Recurrent or Advanced Mismatch Repair–Deficient Endometrial Cancer. *JAMA Oncology.* 2020;6(11):1766. doi:https://doi.org/10.1001/jamaoncol.2020.4515
- 192e. Cacciotti C, Choi J, Alexandrescu S, et al. Immune checkpoint inhibition for pediatric patients with recurrent/refractory CNS tumors: a single institution experience. *Journal of Neuro-Oncology.* 2020;149(1):113-122. doi:https://doi.org/10.1007/s11060-020-03578-6
- 193e. Chalabi M, Verschoor YL, Pedro Batista Tan, et al. Neoadjuvant Immunotherapy in Locally Advanced Mismatch Repair–Deficient Colon Cancer. *New England journal of medicine/The New England journal of medicine.* 2024;390(21):1949-1958. doi:https://doi.org/10.1056/nejmoa2400634
- 194e. Zhang J, Cai J, Deng Y, Wang H. Complete response in patients with locally advanced rectal cancer after neoadjuvant treatment with nivolumab. *Oncolimmunology.* 2019;8(12):e1663108. doi:https://doi.org/10.1080/2162402x.2019.1663108
- 195e. Cercek A, Lumish M, Sinopoli J, et al. PD-1 Blockade in Mismatch Repair–Deficient, Locally Advanced Rectal Cancer. *New England Journal of Medicine.* 2022;386(25). doi:https://doi.org/10.1056/nejmoa2201445
- 196e. Zinzani PL, Ribrag V, Moskowitz CH, et al. Safety and tolerability of pembrolizumab in patients with relapsed/refractory primary mediastinal large B-cell lymphoma. *Blood.* 2017 Jul 20;130(3):267-270. Doi: 10.1182/blood-2016-12-758383. Epub 2017 May 10.

- 197e. Armand P, Rodig S, Melnichenko V, et al. Pembrolizumab in Relapsed or Refractory Primary Mediastinal Large B-Cell Lymphoma. *J Clin Oncol*. 2019 Dec 1;37(34):3291-3299. Doi: 10.1200/JCO.19.01389. Epub 2019 Oct 14
- 198e. You B, Bolze P, Lotz J, et al. Avelumab in Patients With Gestational Trophoblastic Tumors With Resistance to Single-Agent Chemotherapy: Cohort A of the TROPHIMMUN Phase II Trial. *J Clin Oncol*. 2020;38(27):3129-3137. DOI: 10.1200/JCO.20.00803
- 199e. Rudin CM, Awad MM, Navarro A, et al. Pembrolizumab or Placebo Plus Etoposide and Platinum as First-Line Therapy for Extensive-Stage Small-Cell Lung Cancer: Randomized, Double-Blind, Phase III KEYNOTE-604 Study. *J Clin Oncol*. 2020;38(21):2369-2379. Doi:10.1200/JCO.20.00793
- 200e. O'Malley DM, Bariani GM, Cassier PA, et al. Pembrolizumab in Patients With Microsatellite Instability–High Advanced Endometrial Cancer: Results From the KEYNOTE-158 Study. *Journal of Clinical Oncology*. 2022;40(7):752-761. doi:https://doi.org/10.1200/jco.21.01874
- 201e. Santin AD, Deng W, Frumovitz M, et al. Phase II evaluation of nivolumab in the treatment of persistent or recurrent cervical cancer (NCT02257528/NRG-GY002). *Gynecologic Oncology*. 2020;157(1):161-166. doi:https://doi.org/10.1016/j.ygyno.2019.12.034
- 202e. Srinivasan R, Sandeep Gurram, Munjid Al Harthy, et al. Results from a phase II study of bevacizumab and erlotinib in subjects with advanced hereditary leiomyomatosis and renal cell cancer (HLRCC) or sporadic papillary renal cell cancer. *Journal of Clinical Oncology*. 2020;38(15_suppl):5004-5004. doi:https://doi.org/10.1200/jco.2020.38.15_suppl.5004
- 203e. Wakelee H, Liberman M, Kato T, et al. Perioperative Pembrolizumab for Early-Stage Non–Small-Cell Lung Cancer. Published online June 3, 2023. doi:https://doi.org/10.1056/nejmoa2302983
- 204e. Heymach JV, Harpole DH, Tetsuya Mitsudomi, et al. Perioperative Durvalumab for Resectable Non–Small-Cell Lung Cancer. *The New England Journal of Medicine*. Published online October 23, 2023. doi:https://doi.org/10.1056/nejmoa2304875
- 205e. Shitara K, Ajani JA, Moehler M, et al. Nivolumab plus chemotherapy or ipilimumab in gastro-oesophageal cancer. *Nature*. 2022;603(7903):942-948. doi:https://doi.org/10.1038/s41586-022-04508-4
- 206e. Liu T, Jin B, Chen J, et al. Comparative risk of serious and fatal treatment-related adverse events caused by 19 immune checkpoint inhibitors used in cancer treatment: a network meta-analysis. *Therapeutic Advances in Medical Oncology*. 2020;12:175883592094092. doi:https://doi.org/10.1177/1758835920940927
- 207e. P. Borchmann, Moccia AA, Greil R, et al. BRECADDS IS NON-INFERIOR TO EBEACOPP IN PATIENTS WITH ADVANCED STAGE CLASSICAL HODGKIN LYMPHOMA: EFFICACY RESULTS OF THE GHSG PHASE III HD21 TRIAL. *Hematological Oncology*. 2023;41(S2):881-882. doi:https://doi.org/10.1002/hon.3196_lba5

- 208e. Merli F, Stefano Luminari, Gobbi PG, et al. Long-Term Results of the HD2000 Trial Comparing ABVD Versus BEACOPP Versus COPP-EBV-CAD in Untreated Patients With Advanced Hodgkin Lymphoma: A Study by Fondazione Italiana Linfomi. *Journal of Clinical Oncology*. 2015;34(11):1175-1181. doi:<https://doi.org/10.1200/jco.2015.62.4817>
- 209e. Gordon LI, Hong F, Fisher RI, et al. Randomized phase III trial of ABVD versus Stanford V with or without radiation therapy in locally extensive and advanced-stage Hodgkin lymphoma: an intergroup study coordinated by the Eastern Cooperative Oncology Group (E2496). *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2013;31(6):684-691. doi:<https://doi.org/10.1200/JCO.2012.43.4803>
- 210e. Ansell SM, Radford J, Connors JM, et al. Overall Survival with Brentuximab Vedotin in Stage III or IV Hodgkin's Lymphoma. *New England Journal of Medicine*. 2022;387(4):310-320. doi:<https://doi.org/10.1056/nejmoa2206125>
- 211e. Li J, Xu Y, Zang A, et al. Tislelizumab in previously treated, locally advanced unresectable/metastatic microsatellite instability-high/mismatch repair-deficient solid tumors. *PubMed*. 2024;36(3):257-269. doi:<https://doi.org/10.21147/j.issn.1000-9604.2024.03.03>
- 212e. Migden MR, Khushalani NI, Chang ALS, et al. Cemiplimab in locally advanced cutaneous squamous cell carcinoma: results from an open-label, phase 2, single-arm trial. *The Lancet Oncology*. 2020;21(2):294-305. doi:[https://doi.org/10.1016/s1470-2045\(19\)30728-4](https://doi.org/10.1016/s1470-2045(19)30728-4)
- 213e. André T, Berton D, Curigliano G, et al. Antitumor Activity and Safety of Dostarlimab Monotherapy in Patients With Mismatch Repair Deficient Solid Tumors: A Nonrandomized Controlled Trial. *JAMA network open*. 2023;6(11):e2341165. doi:<https://doi.org/10.1001/jamanetworkopen.2023.41165>
- 214e. Chen Y, Wang Y, Zhang H, et al. Short-course radiotherapy combined with chemotherapy and PD-1 inhibitor in low-lying early rectal cancer: study protocol for a single-arm, multicentre, prospective, phase II trial (TORCH-E). *BMJ Open*. 2023;13(10):e076048-e076048. doi:<https://doi.org/10.1136/bmjopen-2023-076048>
- 215e. Lakhani N, Cosman R, Banerji U, et al. A first-in-human phase I study of the PD-1 inhibitor, retifanlimab (INCMGA00012), in patients with advanced solid tumors (POD1UM-101). *ESMO Open*. 2024;9(4):102254-102254. doi:<https://doi.org/10.1016/j.esmoop.2024.102254>
- 216e. Jain N, Jayastu Senapati, Beenu Thakral, et al. A phase 2 study of nivolumab combined with ibrutinib in patients with diffuse large B-cell Richter transformation of CLL. *Blood Advances*. 2023;7(10):1958-1966. doi:<https://doi.org/10.1182/bloodadvances.2022008790>
- 217e. Armand P, Murawski N, Molin D, et al. Pembrolizumab in relapsed or refractory Richter syndrome. *British Journal of Haematology*. 2020;190(2). doi:<https://doi.org/10.1111/bjh.16762>
- 218e. Ding W, LaPlant BR, Call TG, et al. Pembrolizumab in patients with CLL and Richter transformation or with relapsed CLL. *Blood*. 2017;129(26):3419-3427. doi:<https://doi.org/10.1182/blood-2017-02-765685>

- 219e. Othman Al-Sawaf, Ligtoet R, Robrecht S, et al. Tislelizumab plus zanubrutinib for Richter transformation: the phase 2 RT1 trial. *Nature Medicine*. Published online December 9, 2023. doi:<https://doi.org/10.1038/s41591-023-02722-9>
- 220e. Quincy Siu-Chung Chu, Perrone F, L. Greillier, et al. Pembrolizumab plus chemotherapy versus chemotherapy in untreated advanced pleural mesothelioma in Canada, Italy, and France: a phase 3, open-label, randomised controlled trial. *The Lancet*. Published online November 1, 2023. doi:[https://doi.org/10.1016/s0140-6736\(23\)01613-6](https://doi.org/10.1016/s0140-6736(23)01613-6)
- 221e. Zalcman G, Mazieres J, Margery J, et al. Bevacizumab for newly diagnosed pleural mesothelioma in the Mesothelioma Avastin Cisplatin Pemetrexed Study (MAPS): a randomised, controlled, open-label, phase 3 trial. *The Lancet*. 2016;387(10026):1405-1414. doi:[https://doi.org/10.1016/S0140-6736\(15\)01238-6](https://doi.org/10.1016/S0140-6736(15)01238-6)
- 222e. Krug LM, Pass HI, Rusch VW, et al. Multicenter Phase II Trial of Neoadjuvant Pemetrexed Plus Cisplatin Followed by Extrapleural Pneumonectomy and Radiation for Malignant Pleural Mesothelioma. *Journal of Clinical Oncology*. 2009;27(18):3007-3013. doi:<https://doi.org/10.1200/jco.2008.20.3943>
- 223e. Thieke C, Nicolay NH, Florian Sterzing, et al. Long-term results in malignant pleural mesothelioma treated with neoadjuvant chemotherapy, extrapleural pneumonectomy and intensity-modulated radiotherapy. 2015;10(1). doi:<https://doi.org/10.1186/s13014-015-0575-5>
- 224e. Ignace Vergote, Els Van Nieuwenhuysen, O'Cearbhaill RE, et al. Tisotumab Vedotin in Combination With Carboplatin, Pembrolizumab, or Bevacizumab in Recurrent or Metastatic Cervical Cancer: Results From the innovaTV 205/GOG-3024/ENGOT-cx8 Study. *Journal of Clinical Oncology*. 2023;41(36):5536-5549. doi:<https://doi.org/10.1200/jco.23.00720>
- 225e. Seok Hyun Kim, Jing Quan Lim, Yuriike Laurensia, et al. Avelumab for the treatment of relapsed or refractory extranodal NK/T-cell lymphoma: an open-label phase 2 study. *Blood*. 2020;136(24):2754-2763. doi:<https://doi.org/10.1182/blood.2020007247>
- 226e. Patel SP, Othus M, Chae YK, et al. A Phase II Basket Trial of Dual Anti-CTLA-4 and Anti-PD-1 Blockade in Rare Tumors (DART SWOG 1609 Cohort 47) in Patients with Gestational Trophoblastic Neoplasia. *Clinical Cancer Research*. 2023;30(1):33-38. doi:<https://doi.org/10.1158/1078-0432.ccr-23-2293>
- 227e. Delyon J, Bizot A, Battistella M, Madelaine I, Vercellino L, Lebbé C. PD-1 blockade with nivolumab in endemic Kaposi sarcoma. *Annals of Oncology*. 2018;29(4):1067-1069. doi:<https://doi.org/10.1093/annonc/mdy006>
- 228e. Florou V, Rosenberg AE, Wieder E, et al. Angiosarcoma patients treated with immune checkpoint inhibitors: a case series of seven patients from a single institution. *Journal for ImmunoTherapy of Cancer*. 2019;7(1). doi:<https://doi.org/10.1186/s40425-019-0689-7>
- 229e. Maki RG, Wathen JK, Patel SR, et al. Randomized Phase II Study of Gemcitabine and Docetaxel Compared With Gemcitabine Alone in Patients With Metastatic Soft Tissue

- Sarcomas: Results of Sarcoma Alliance for Research Through Collaboration Study 002. *Journal of Clinical Oncology*. 2007;25(19):2755-2763. doi:<https://doi.org/10.1200/jco.2006.10.4117>
- 230e. Samuels BL, Chawla SP, Patel S, et al. Clinical outcomes and safety with trabectedin therapy in patients with advanced soft tissue sarcomas following failure of prior chemotherapy: results of a worldwide expanded access program study. *Annals of Oncology*. 2013;24(6):1703-1709. doi:<https://doi.org/10.1093/annonc/mds659>
- 231e. García-Del-Muro X, López-Pousa A, Maurel J, et al. Randomized phase II study comparing gemcitabine plus dacarbazine versus dacarbazine alone in patients with previously treated soft tissue sarcoma: a Spanish Group for Research on Sarcomas study. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*. 2011;29(18):2528-2533. doi:<https://doi.org/10.1200/JCO.2010.33.6107>
- 232e. D'Angelo SP, Mahoney MR, Van Tine BA, et al. Nivolumab with or without ipilimumab treatment for metastatic sarcoma (Alliance A091401): two open-label, non-comparative, randomised, phase 2 trials. *The Lancet Oncology*. 2018;19(3):416-426. doi:[https://doi.org/10.1016/S1470-2045\(18\)30006-8](https://doi.org/10.1016/S1470-2045(18)30006-8)
- 233e. Melissa Amber Burgess, Bolejack V, Van BA, et al. Multicenter phase II study of pembrolizumab (P) in advanced soft tissue (STS) and bone sarcomas (BS): Final results of SARC028 and biomarker analyses. 2017;35(15_suppl):11008-11008. doi:https://doi.org/10.1200/jco.2017.35.15_suppl.11008
- 234e. Judson I, Verweij J, Gelderblom H, et al. Doxorubicin alone versus intensified doxorubicin plus ifosfamide for first-line treatment of advanced or metastatic soft-tissue sarcoma: a randomised controlled phase 3 trial. *The Lancet Oncology*. 2014;15(4):415-423. doi:[https://doi.org/10.1016/s1470-2045\(14\)70063-4](https://doi.org/10.1016/s1470-2045(14)70063-4)
- 235e. van der Graaf WT, Blay JY, Chawla SP, et al. Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial. *The Lancet*. 2012;379(9829):1879-1886. doi:[https://doi.org/10.1016/s0140-6736\(12\)60651-5](https://doi.org/10.1016/s0140-6736(12)60651-5)
- 236e. Schöffski P, Ray-Coquard I, Cioffi A, et al. Activity of eribulin mesylate in patients with soft-tissue sarcoma: a phase 2 study in four independent histological subtypes. 2011;12(11):1045-1052. doi:[https://doi.org/10.1016/s1470-2045\(11\)70230-3](https://doi.org/10.1016/s1470-2045(11)70230-3)
- 237e. Agulnik M, Schulte B, Robinson S, et al. An open-label single-arm phase II study of regorafenib for the treatment of angiosarcoma. *European Journal of Cancer*. 2021;154:201-208. doi:<https://doi.org/10.1016/j.ejca.2021.06.027>
- 238e. Penel N, Bui B, Bay JO, et al. Phase II Trial of Weekly Paclitaxel for Unresectable Angiosarcoma: The ANGIOTAX Study. *Journal of Clinical Oncology*. 2008;26(32):5269-5274. doi:<https://doi.org/10.1200/jco.2008.17.3146>
- 239e. Verweij J, Lee SM, Ruka W, et al. Randomized Phase II Study of Docetaxel Versus Doxorubicin in First- and Second-Line Chemotherapy for Locally Advanced or Metastatic Soft

Tissue Sarcomas in Adults: A Study of the European Organization for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group. *Journal of Clinical Oncology*. 2000;18(10):2081-2086. doi:<https://doi.org/10.1200/jco.2000.18.10.2081>

- 240e. Blank CU, et al. Neoadjuvant Nivolumab and Ipilimumab in Resectable Stage III Melanoma. *New England journal of medicine/The New England journal of medicine*. Published online June 2, 2024. doi:<https://doi.org/10.1056/nejmoa2402604>
- 241e. Rosa K. Lenvatinib/Pembrolizumab Combo Proves Safe, Effective for Metastasized Anaplastic and Poorly Differentiated Thyroid Carcinoma. OncLive. Published September 13, 2022. Accessed January 21, 2025. <https://www.onclive.com/view/lenvatinib-pembrolizumab-combo-proves-safe-effective-for-metastasized-anaplastic-and-poorly-differentiated-thyroid-carcinoma>
- 242e. Ain KB, Egorin MJ, DeSimone PA. Treatment of Anaplastic Thyroid Carcinoma with Paclitaxel: Phase 2 Trial Using Ninety-Six-hour Infusion. *Thyroid*. 2000;10(7):587-594. doi:<https://doi.org/10.1089/thy.2000.10.587>
- 243e. Shimaoka K, Schoenfeld DA, Dewys WD, Creech RH, Deconti R. A randomized trial of doxorubicin versus doxorubicin plus cisplatin in patients with advanced thyroid carcinoma. *Cancer*. 1985;56(9):2155-2160. doi:[https://doi.org/10.1002/1097-0142\(19851101\)56:9%3C2155::aid-cnrc2820560903%3E3.0.co;2-e](https://doi.org/10.1002/1097-0142(19851101)56:9%3C2155::aid-cnrc2820560903%3E3.0.co;2-e)
- 244e. Oh DY, He AR, Qin S, et al. A phase 3 randomized, double-blind, placebo-controlled study of durvalumab in combination with gemcitabine plus cisplatin (GemCis) in patients (pts) with advanced biliary tract cancer (BTC): TOPAZ-1. *J Clin Oncol*. 2022 Feb 1;40(4_suppl):378-378.
- 245e. Hoimes CJ, Flaig TW, Milowsky MI, et al. Enfortumab Vedotin Plus Pembrolizumab in Previously Untreated Advanced Urothelial Cancer. *J Clin Oncol*. 2023 Jan 1;41(1):22-31. doi: 10.1200/JCO.22.01643.
- 246e. Apolo AB, Ballman KV, Sonpavde G, et al. Adjuvant Pembrolizumab versus Observation in Muscle-Invasive Urothelial Carcinoma. *N Engl J Med*. 2025;392(1):45-55
- 247e. Coleman RL, Lorusso D, Gennigens C, et al; innovaTV 204/GOG-3023/ENGOT-cx6 Collaborators. Efficacy and safety of tisotumab vedotin in previously treated recurrent or metastatic cervical cancer (innovaTV 204/GOG-3023/ENGOT-cx6): a multicentre, open-label, single-arm, phase 2 study. *Lancet Oncol*. 2021 May;22(5):609-619.
- 248e. Georger B, Kang HJ, Yalon-Oren M. Pembrolizumab in paediatric patients with advanced melanoma or a PD-L1-positive, advanced, relapsed, or refractory solid tumour or lymphoma (KEYNOTE-051): interim analysis of an open-label, single-arm, phase 1-2 trial. *Lancet Oncol*. 2020 Jan;21(1):121-133. doi: 10.1016/S1470-2045(19)30671-0. Epub 2019 Dec 4. PMID: 31812554.
- 249e. Ferris RL, Blumenschein G Jr, Fayette J, et al. Nivolumab vs investigator's choice in recurrent or metastatic squamous cell carcinoma of the head and neck: 2-year long-term survival update

- of CheckMate 141 with analyses by tumor PD-L1 expression. *Oral Oncol.* 2018 Jun;81:45-51. doi: 10.1016/j.oraloncology.2018.04.008. Epub 2018 Apr 17.
- 250e. Friedberg JW, Forero-Torres A, Bordoni RE, et al. Frontline brentuximab vedotin in combination with dacarbazine or bendamustine in patients aged ≥ 60 years with HL. *Blood.* 2017 Dec 28;130(26):2829-2837.
- 251e. Abou-Alfa GK, Chan SL, Kudo M, et al. Phase 3 randomized, open-label, multicenter study of tremelimumab (T) and durvalumab (D) as first-line therapy in patients (pts) with unresectable hepatocellular carcinoma (uHCC): HIMALAYA. *Journal of Clinical Oncology* 2022 40:4_suppl, 379-379.
- 252e. Abou-Alfa GK, Chan SL, Furuse J, et al. A randomized, multicenter phase 3 study of durvalumab (D) and tremelimumab (T) as first-line treatment in patients with unresectable hepatocellular carcinoma (HCC): HIMALAYA study. *Journal of Clinical Oncology* 36, no. 15_suppl. DOI: 10.1200/JCO.2018.36.15_suppl.TPS4144.
- 253e. Llovet JM, Ricci S, Mazzaferro V; SHARP Investigators Study Group. Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med.* 2008 Jul 24;359(4):378-90. doi: 10.1056/NEJMoa0708857. PMID: 18650514.
- 254e. Kudo M, Finn RS, Qin S. Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. *Lancet.* 2018 Mar 24;391(10126):1163-1173. doi: 10.1016/S0140-6736(18)30207-1. PMID: 29433850.
- 255e. Finn RS, Qin S, Ikeda M, et al; IMbrave150 Investigators. Atezolizumab plus Bevacizumab in Unresectable Hepatocellular Carcinoma. *N Engl J Med.* 2020 May 14;382(20):1894-1905.
- 256e. Trigo J, Subbiah V, Besse B, et al. Lurbinectedin as second-line treatment for patients with small-cell lung cancer: a single-arm, open-label, phase 2 basket trial. *Lancet Oncol.* 2020 May;21(5):645-654. doi: 10.1016/S1470-2045(20)30068-1. Epub 2020 Mar 27.
- 257e. Eckardt JR, von Pawel J, Pujol JL, et al. Phase III study of oral compared with intravenous topotecan as second-line therapy in small-cell lung cancer. *J Clin Oncol.* 2007 May 20;25(15):2086-92.
- 258e. Edelman MJ, Dvorkin M, Laktionov K, et al. Randomized phase 3 study of the anti-disialoganglioside antibody dinutuximab and irinotecan vs irinotecan or topotecan for second-line treatment of small cell lung cancer. *Lung Cancer.* 2022;166:135-142
- 259e. Baize N, Monnet I, Greillier L, et al. Carboplatin plus etoposide versus topotecan as second-line treatment for patients with sensitive relapsed small-cell lung cancer: an open-label, multicentre, randomised, phase 3 trial. *Lancet Oncol* 2020;21:1224-1233.
- 260e. Naito Y, Yamada K, Imamura Y, et al. Rechallenge treatment with a platinum-based regimen in patients with sensitive relapsed small-cell lung cancer. *Med Oncol* 2018;35:61.

- 261e. Genestreti G, Tiseo M, Kenmotsu H, et al. Outcomes of platinum-sensitive small-cell lung cancer patients treated with platinum/etoposide rechallenge: a multi-institutional retrospective analysis. *Clin Lung Cancer* 2015;16:e223-e228
- 262e. Rousseau B, Bièche I, Éric Pasmant, et al. PD-1 Blockade in Solid Tumors with Defects in Polymerase Epsilon. *Cancer Discovery*. 2022;12(6):1435-1448. doi:<https://doi.org/10.1158/2159-8290.cd-21-0521>
- 263e. Janjigian YY, Ajani JA, Moehler M, et al. First-Line Nivolumab Plus Chemotherapy for Advanced Gastric, Gastroesophageal Junction, and Esophageal Adenocarcinoma: 3-Year Follow-Up of the Phase III CheckMate 649 Trial. *J Clin Oncol* 2024 Feb 21;42(17):2012–2020.
- 264e. Mei MG, Lee HJ, Palmer JM, et al. Response-adapted anti-PD-1-based salvage therapy for Hodgkin lymphoma with nivolumab alone or in combination with ICE. *Blood* 2022;139:3605-3616
- 265e. Bryan LJ, Casulo C, Allen PB, et al. Pembrolizumab Added to Ifosfamide, Carboplatin, and Etoposide Chemotherapy for Relapsed or Refractory Classic Hodgkin Lymphoma: A Multi-institutional Phase 2 Investigator-Initiated Nonrandomized Clinical Trial. *JAMA Oncol*. 2023 May 1;9(5):683-691. doi: 10.1001/jamaoncol.2022.7975. PMID: 36928527; PMCID: PMC10020934.
- 266e. Raj N, Zheng Y, Kelly V, et al. PD-1 Blockade in Advanced Adrenocortical Carcinoma. *J Clin Oncol*. 2020 Jan 1;38(1):71-80. Doi: 10.1200/JCO.19.01586.
- 267e. Naing A, Meric-Bernstam F, Stephen B, et al. Phase 2 study of pembrolizumab in patients with advanced rare cancers [published correction appears in *J Immunother Cancer*. 2020 Apr;8(1):]. *J Immunother Cancer*. 2020;8(1):e000347. Doi:10.1136/jitc-2019-000347.
- 268e. Munhoz RR, Nader-Marta G, de Camargo VP, et al. A phase 2 study of first-line nivolumab in patients with locally advanced or metastatic cutaneous squamous-cell carcinoma. *Cancer* 2022;128:4223- 4231.
- 269e. Duan H, Shao C, Pan M, et al. Neoadjuvant Pembrolizumab and Chemotherapy in Resectable Esophageal Cancer: An Open-Label, Single-Arm Study (PEN-ICE). *Front Immunol*. 2022 Jun 2;13:849984. doi: 10.3389/fimmu.2022.849984. PMID: 35720388; PMCID: PMC9202755.
- 270e. Das A, Tabori U, Sambira Nahum L, et al. Efficacy of Nivolumab in Pediatric Cancers with High Mutation Burden and Mismatch Repair Deficiency. *Clin Cancer Res*. 2023 Dec 1;29(23):4770-4783. doi: 10.1158/1078-0432.CCR-23-0411.
- 271e. Das S, Allen A, Berlin J. Immunotherapy After Immunotherapy: Response Rescue in a Patient With Microsatellite Instability-high Colorectal Cancer Post-Pembrolizumab. *Clin Colorectal Cancer*. 2020 Jun;19(2):137-140. doi: 10.1016/j.clcc.2020.02.006.
- 272e. Klein O, Senko C, Carlino MS, et al. Combination immunotherapy with ipilimumab and nivolumab in patients with advanced adrenocortical carcinoma: a subgroup analysis of CA209-538. *Oncoimmunology*. 2021 Apr 12;10(1):1908771. doi: 10.1080/2162402X.2021.1908771.

- 273e. Bhatia S, Topalian S, Sharfman W, et al. Nivolumab With or Without Ipilimumab in Patients With Recurrent or Metastatic Merkel Cell Carcinoma: A Nonrandomized, Open-Label, International, Multicenter Phase I/II Study. *J Clin Oncol*. 2025 Jan 31;43(9):1137–1147. doi: 10.1200/JCO-24-02138
- 274e. Paoluzzi L, Cacavio A, Ghesani M, et al. Response to anti-PD1 therapy with nivolumab in metastatic sarcomas. *Clin Sarcoma Res*. 2016 Dec 30;6:24. doi: 10.1186/s13569-016-0064-0. eCollection 2016.
- 275e. Tawbi HA, Burgess M, Bolejack V, et al. Pembrolizumab in advanced soft-tissue sarcoma and bone sarcoma (SARC028): a multicentre, two-cohort, single-arm, open-label, phase 2 trial. *Lancet Oncol*. 2017 Nov;18(11):1493-1501. doi: 10.1016/S1470-2045(17)30624-1.
- 276e. Schenker M, Burotto M, Richardet M, et al. Randomized, open-label, phase 2 study of nivolumab plus ipilimumab or nivolumab monotherapy in patients with advanced or metastatic solid tumors of high tumor mutational burden. *J Immunother Cancer*. 2024 Aug 6;12(8):e008872. doi: 10.1136/jitc-2024-008872.
- 277e. Tewari KS, Monk BJ, Vergote I, et al. Survival with Cemiplimab in Recurrent Cervical Cancer. *N Engl J Med* 2022;386:544-555.
- 278e. Cohen R, Palmieri LJ, Boisson C et al. Nivolumab and ipilimumab in anti-PD1-resistant MSI/dMMR metastatic colorectal cancer: Interim analysis of the GERCOR NIPIRESCUE phase II trial. *Ann Oncol*. 2025;36(Suppl2):S545. Abstract 879P.
- 279e. Cercek A, Foote MB, Rousseau B, et al. Non-operative management of mismatch repair-deficient tumors. *N Engl J Med* 2025; 392:2297-2308.
- 280e. Chanza NM, Xie E, Bilen MA, et al. Cabozantinib in advanced non-clear-cell renal cell carcinoma: a multicentre, retrospective, cohort study. *Lancet Oncol*. 2019 Apr;20(4):581-590. doi: 10.1016/S1470-2045(18)30907-0.
- 281e. André T, Elez E, Lenz H et al. Nivolumab plus ipilimumab versus nivolumab in microsatellite instability-high metastatic colorectal cancer (CheckMate 8HW): a randomised, open-label, phase 3 trial. *The Lancet*, 2025; 405, 383-395
- 282e. Morris VK, Ciombor KK, Xiao L, Ochieng JK, Marmonti E, Polite B, Weinberg BA, Krauss JC, Hays J, Mukherjee S, Aranha O, Iqbal S, Shields T, Benson AB, Kazmi S, Lieu C, Hochster H, Whisenant J, Haymaker C, Eng C. NCI9673 (Part B): ETCTN Randomized Phase II Study of Nivolumab With or Without Ipilimumab in Refractory, Metastatic Squamous Cell Carcinoma of the Anal Canal. *J Clin Oncol*. 2026 Feb 20;44(6):497-507. doi: 10.1200/JCO-25-00929. Epub 2026 Jan 7. PMID: 41499716; PMCID: PMC12782278.
- 283e. Rao S, Samalin-Scalzi E, Evesque L, et al. LBA2 POD1UM-303/InterAACT 2: Phase III study of retifanlimab with carboplatin-paclitaxel (c-p) in patients (Pts) with inoperable locally recurrent or metastatic squamous cell carcinoma of the anal canal (SCAC) not previously treated with systemic chemotherapy (Chemo)/ *Annals of Oncology*, 35S1217
- 284e. Long GV, Atkinson V, Lo SN, et al. Ipilimumab plus nivolumab versus nivolumab alone in patients with melanoma brain metastases (ABC): 7-year follow-up of a multicentre, open-label, randomised, phase 2 study. *Lancet Oncol*. 2025 Mar;26(3):320-330. doi: 10.1016/S1470-2045(24)00735-6. Epub 2025 Feb 17. PMID: 39978375.

- 285e. Rocha Lima, Yothers G, George TJ, et al. Colorectal cancer metastatic dMMR immunotherapy (COMMIT) study: a randomized phase III study of atezolizumab (atezo) monotherapy versus mFOLFOX6/bevacizumab/atezo (FFX/bev) in the first-line treatment of patients (pts) with deficient DNA mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer (mCRC)— NRG-GI004/SWOG-S1610. *J Clin Oncol*. 2026;44(Suppl 2):14. doi:10.1200/JCO.2026.44.2_suppl.14
- 286e. Friedman CF, Hainsworth JD, et al. Atezolizumab Treatment of Tumors with High Tumor Mutational Burden from MyPathway, a Multicenter, Open-Label, Phase IIa Multiple Basket Study. *Cancer Discov*. 2022 Mar 1;12(3):654-669. doi: 10.1158/2159-8290.CD-21-0450. PMID: 34876409; PMCID: PMC9394388.
- 287e. Chen G, Ye Y, Liu Y, Sun Y, Gao Y, et al. Efficacy and safety of tislelizumab as neoadjuvant treatment in patients with stage II-III microsatellite instability-high/mismatch repair-deficient colorectal cancer: a single-arm, multicenter, open-label, prospective, phase II study. *Int J Surg*. 2026 Feb 18. doi: 10.1097/JS9.0000000000004836. Epub ahead of print. PMID: 41706635.
- 288e. Burtneß B, Harrington KJ, Greil R, Soulières D, et al.; KEYNOTE-048 Investigators. Pembrolizumab alone or with chemotherapy versus cetuximab with chemotherapy for recurrent or metastatic squamous cell carcinoma of the head and neck (KEYNOTE-048): a randomised, open-label, phase 3 study. *Lancet*. 2019 Nov 23;394(10212):1915-1928. doi: 10.1016/S0140-6736(19)32591-7. Epub 2019 Nov 1.
- 289e. Rimassa L, Chan SL, Sangro B, Lau G, et al. Five-year overall survival update from the HIMALAYA study of tremelimumab plus durvalumab in unresectable HCC. *J Hepatol*. 2025 Oct;83(4):899-908. doi: 10.1016/j.jhep.2025.03.033. Epub 2025 Apr 11. PMID: 40222621.
- 290e. Melero I, Yau T, Kang YK, Kim TY, Santoro A, et al. Nivolumab plus ipilimumab combination therapy in patients with advanced hepatocellular carcinoma previously treated with sorafenib: 5-year results from CheckMate 040. *Ann Oncol*. 2024 Jun;35(6):537-548. doi: 10.1016/j.annonc.2024.03.005. Epub 2024 May 22. PMID: 38844309.
- 291e. Kim HD, Sym SJ, Chon HJ, et al. Multicenter single-arm phase II trial of lenvatinib in patients with advanced hepatocellular carcinoma after progression on first-line atezolizumab plus bevacizumab. *J Hepatol*. 2026 Feb;84(2):308-315. doi: 10.1016/j.jhep.2025.08.020. Epub 2025 Sep 4. PMID: 41038766.
- 292e. Friedman CF, Manning-Geist BL, Zhou Q, et al. Nivolumab for mismatch-repair-deficient or hypermutated gynecologic cancers: a phase 2 trial with biomarker analyses. *Nat Med*. 2024 May;30(5):1330-1338. doi: 10.1038/s41591-024-02942-7. Epub 2024 Apr 23. PMID: 38653864; PMCID: PMC11108776.
- 293e. Dizon DS, Mathews CA, David SM, et al. Final results of BrUOG 354: A randomized phase II trial of nivolumab alone or in combination with ipilimumab for people with ovarian and other extra-renal clear cell carcinomas.. *J Clin Oncol* 42, LBA5500-LBA5500(2024). DOI:10.1200/JCO.2024.42.17_suppl.LBA5500
- 294e. Prime Therapeutics Management. Opdivo Clinical Literature Review Analysis. Last updated April 2026. Accessed April 2026.

Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime's assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	No: PA not a priority
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C00.3	Malignant neoplasm of upper lip, inner aspect
C00.4	Malignant neoplasm of lower lip, inner aspect
C00.5	Malignant neoplasm of lip, unspecified, inner aspect
C00.6	Malignant neoplasm of commissure of lip, unspecified
C00.8	Malignant neoplasm of overlapping sites of lip
C00.9	Malignant neoplasm of lip, unspecified
C01	Malignant neoplasm of base of tongue
C02.0	Malignant neoplasm of dorsal surface of tongue
C02.1	Malignant neoplasm of border of tongue
C02.2	Malignant neoplasm of ventral surface of tongue
C02.3	Malignant neoplasm of anterior two-thirds of tongue, part unspecified
C02.4	Malignant neoplasm of lingual tonsil
C02.8	Malignant neoplasm of overlapping sites of tongue
C02.9	Malignant neoplasm of tongue, unspecified
C03.0	Malignant neoplasm of upper gum
C03.1	Malignant neoplasm of lower gum
C03.9	Malignant neoplasm of gum, unspecified
C04.0	Malignant neoplasm of anterior floor of mouth

Medical Necessity Criteria

C04.1	Malignant neoplasm of lateral floor of mouth
C04.8	Malignant neoplasm of overlapping sites of floor of mouth
C04.9	Malignant neoplasm of floor of mouth, unspecified
C05.0	Malignant neoplasm of hard palate
C05.1	Malignant neoplasm of soft palate
C05.2	Malignant neoplasm of uvula
C05.8	Malignant neoplasm of overlapping sites of palate
C05.9	Malignant neoplasm of palate, unspecified
C06.0	Malignant neoplasm of cheek mucosa
C06.2	Malignant neoplasm of retromolar area
C06.80	Malignant neoplasm of overlapping sites of unspecified parts of mouth
C06.89	Malignant neoplasm of overlapping sites of other parts of mouth
C06.9	Malignant neoplasm of mouth, unspecified
C09.0	Malignant neoplasm of tonsillar fossa
C09.1	Malignant neoplasm of tonsillar pillar (anterior) (posterior)
C09.8	Malignant neoplasm of overlapping sites of tonsil
C09.9	Malignant neoplasm of tonsil, unspecified
C10.0	Malignant neoplasm of vallecula
C10.1	Malignant neoplasm of anterior surface of epiglottis
C10.2	Malignant neoplasm of lateral wall of oropharynx
C10.3	Malignant neoplasm of posterior wall of oropharynx
C10.4	Malignant neoplasm of branchial cleft
C10.8	Malignant neoplasm of overlapping sites of oropharynx
C10.9	Malignant neoplasm of oropharynx, unspecified
C12	Malignant neoplasm of pyriform sinus
C13.0	Malignant neoplasm of postcricoid region
C13.1	Malignant neoplasm of aryepiglottic fold, hypopharyngeal aspect
C13.2	Malignant neoplasm of posterior wall of hypopharynx
C13.8	Malignant neoplasm of overlapping sites of hypopharynx
C13.9	Malignant neoplasm of hypopharynx, unspecified
C14.0	Malignant neoplasm of pharynx, unspecified

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C14.2	Malignant neoplasm of Waldeyer's ring
C14.8	Malignant neoplasm of overlapping sites of lip, oral cavity and pharynx
C15.3	Malignant neoplasm of upper third of esophagus
C15.4	Malignant neoplasm of middle third of esophagus
C15.5	Malignant neoplasm of lower third of esophagus
C15.8	Malignant neoplasm of overlapping sites of esophagus
C15.9	Malignant neoplasm of esophagus, unspecified
C16.0	Malignant neoplasm of cardia
C16.1	Malignant neoplasm of fundus of stomach
C16.2	Malignant neoplasm of body of stomach
C16.3	Malignant neoplasm of pyloric antrum
C16.4	Malignant neoplasm of pylorus
C16.5	Malignant neoplasm of lesser curvature of stomach, unspecified
C16.6	Malignant neoplasm of greater curvature of stomach, unspecified
C16.8	Malignant neoplasm of overlapping sites of stomach
C16.9	Malignant neoplasm of stomach, unspecified
C17.0	Malignant neoplasm of duodenum
C17.1	Malignant neoplasm of jejunum
C17.2	Malignant neoplasm of ileum
C17.3	Meckel's diverticulum, malignant
C17.8	Malignant neoplasm of overlapping sites of small intestine
C17.9	Malignant neoplasm of small intestine, unspecified
C18.0	Malignant neoplasm of cecum
C18.1	Malignant neoplasm of appendix
C18.2	Malignant neoplasm of ascending colon
C18.3	Malignant neoplasm of hepatic flexure
C18.4	Malignant neoplasm of transverse colon
C18.5	Malignant neoplasm of splenic flexure
C18.6	Malignant neoplasm of descending colon
C18.7	Malignant neoplasm of sigmoid colon
C18.8	Malignant neoplasm of overlapping sites of colon

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C18.9	Malignant neoplasm of colon, unspecified
C19	Malignant neoplasm of rectosigmoid junction
C20	Malignant neoplasm of rectum
C21.0	Malignant neoplasm of anus, unspecified
C21.1	Malignant neoplasm of anal canal
C21.2	Malignant neoplasm of cloacogenic zone
C21.8	Malignant neoplasm of overlapping sites of rectum, anus and anal canal
C22.0	Liver cell carcinoma
C22.1	Intrahepatic bile duct carcinoma
C22.3	Angiosarcoma of liver
C22.8	Malignant neoplasm of liver, primary, unspecified as to type
C22.9	Malignant neoplasm of liver, not specified as primary or secondary
C23	Malignant neoplasm of gallbladder
C24.0	Malignant neoplasm of extrahepatic bile duct
C24.1	Malignant neoplasm of ampulla of Vater
C24.8	Malignant neoplasm of overlapping sites of biliary tract
C24.9	Malignant neoplasm of biliary tract, unspecified
C31.0	Malignant neoplasm of maxillary sinus
C31.1	Malignant neoplasm of ethmoidal sinus
C32.0	Malignant neoplasm of glottis
C32.1	Malignant neoplasm of supraglottis
C32.2	Malignant neoplasm of subglottis
C32.3	Malignant neoplasm of laryngeal cartilage
C32.8	Malignant neoplasm of overlapping sites of larynx
C32.9	Malignant neoplasm of larynx, unspecified
C33	Malignant neoplasm of trachea
C34.00	Malignant neoplasm of unspecified main bronchus
C34.01	Malignant neoplasm of right main bronchus
C34.02	Malignant neoplasm of left main bronchus
C34.10	Malignant neoplasm of upper lobe, unspecified bronchus or lung
C34.11	Malignant neoplasm of upper lobe, right bronchus or lung

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C34.12	Malignant neoplasm of upper lobe, left bronchus or lung
C34.2	Malignant neoplasm of middle lobe, bronchus or lung
C34.30	Malignant neoplasm of lower lobe, unspecified bronchus or lung
C34.31	Malignant neoplasm of lower lobe, right bronchus or lung
C34.32	Malignant neoplasm of lower lobe, left bronchus or lung
C34.80	Malignant neoplasm of overlapping sites of unspecified bronchus and lung
C34.81	Malignant neoplasm of overlapping sites of right bronchus and lung
C34.82	Malignant neoplasm of overlapping sites of left bronchus and lung
C34.90	Malignant neoplasm of unspecified part of unspecified bronchus or lung
C34.91	Malignant neoplasm of unspecified part of right bronchus or lung
C34.92	Malignant neoplasm of unspecified part of left bronchus or lung
C40.00	Malignant neoplasm of scapula and long bones of unspecified upper limb
C40.01	Malignant neoplasm of scapula and long bones of right upper limb
C40.02	Malignant neoplasm of scapula and long bones of left upper limb
C40.10	Malignant neoplasm of short bones of unspecified upper limb
C40.11	Malignant neoplasm of short bones of right upper limb
C40.12	Malignant neoplasm of short bones of left upper limb
C40.20	Malignant neoplasm of long bones of unspecified lower limb
C40.21	Malignant neoplasm of long bones of right lower limb
C40.22	Malignant neoplasm of long bones of left lower limb
C40.30	Malignant neoplasm of short bones of unspecified lower limb
C40.31	Malignant neoplasm of short bones of right lower limb
C40.32	Malignant neoplasm of short bones of left lower limb
C40.80	Malignant neoplasm of overlapping sites of bone and articular cartilage of unspecified limb
C40.81	Malignant neoplasm of overlapping sites of bone and articular cartilage of right limb
C40.82	Malignant neoplasm of overlapping sites of bone and articular cartilage of left limb
C40.90	Malignant neoplasm of unspecified bones and articular cartilage of unspecified limb
C40.91	Malignant neoplasm of unspecified bones and articular cartilage of right limb
C40.92	Malignant neoplasm of unspecified bones and articular cartilage of left limb
C41.0	Malignant neoplasm of bones of skull and face
C41.1	Malignant neoplasm of mandible

C41.2	Malignant neoplasm of vertebral column
C41.3	Malignant neoplasm of ribs, sternum and clavicle
C41.4	Malignant neoplasm of pelvic bones, sacrum and coccyx
C41.9	Malignant neoplasm of bone and articular cartilage, unspecified
C43.0	Malignant melanoma of lip
C43.10	Malignant melanoma of unspecified eyelid, including canthus
C43.111	Malignant melanoma of right upper eyelid, including canthus
C43.112	Malignant melanoma of right lower eyelid, including canthus
C43.121	Malignant melanoma of left upper eyelid, including canthus
C43.122	Malignant melanoma of left lower eyelid, including canthus
C43.20	Malignant melanoma of unspecified ear and external auricular canal
C43.21	Malignant melanoma of right ear and external auricular canal
C43.22	Malignant melanoma of left ear and external auricular canal
C43.30	Malignant melanoma of unspecified part of face
C43.31	Malignant melanoma of nose
C43.39	Malignant melanoma of other parts of face
C43.4	Malignant melanoma of scalp and neck
C43.51	Malignant melanoma of anal skin
C43.52	Malignant melanoma of skin of breast
C43.59	Malignant melanoma of other part of trunk
C43.60	Malignant melanoma of unspecified upper limb, including shoulder
C43.61	Malignant melanoma of right upper limb, including shoulder
C43.62	Malignant melanoma of left upper limb, including shoulder
C43.70	Malignant melanoma of unspecified lower limb, including hip
C43.71	Malignant melanoma of right lower limb, including hip
C43.72	Malignant melanoma of left lower limb, including hip
C43.8	Malignant melanoma of overlapping sites of skin
C43.9	Malignant melanoma of skin, unspecified
C44.02	Squamous cell carcinoma of skin of lip
C45.0	Mesothelioma of pleura
C45.1	Mesothelioma of peritoneum

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C45.2	Mesothelioma of pericardium
C45.7	Mesothelioma of other sites
C45.9	Mesothelioma, unspecified
C46.0	Kaposi's sarcoma of skin
C46.1	Kaposi's sarcoma of soft tissue
C46.2	Kaposi's sarcoma of palate
C46.3	Kaposi's sarcoma of lymph nodes
C46.4	Kaposi's sarcoma of gastrointestinal sites
C46.50	Kaposi's sarcoma of unspecified lung
C46.51	Kaposi's sarcoma of right lung
C46.52	Kaposi's sarcoma of left lung
C46.7	Kaposi's sarcoma of other sites
C46.9	Kaposi's sarcoma, unspecified
C47.0	Malignant neoplasm of peripheral nerves of head, face and neck
C47.10	Malignant neoplasm of peripheral nerves of unspecified upper limb, including shoulder
C47.11	Malignant neoplasm of peripheral nerves of right upper limb, including shoulder
C47.12	Malignant neoplasm of peripheral nerves of left upper limb, including shoulder
C47.20	Malignant neoplasm of peripheral nerves of unspecified lower limb, including hip
C47.21	Malignant neoplasm of peripheral nerves of right lower limb, including hip
C47.22	Malignant neoplasm of peripheral nerves of left lower limb, including hip
C47.3	Malignant neoplasm of peripheral nerves of thorax
C47.4	Malignant neoplasm of peripheral nerves of abdomen
C47.5	Malignant neoplasm of peripheral nerves of pelvis
C47.6	Malignant neoplasm of peripheral nerves of trunk, unspecified
C47.8	Malignant neoplasm of overlapping sites of peripheral nerves and autonomic nervous system
C47.9	Malignant neoplasm of peripheral nerves and autonomic nervous system, unspecified
C48.0	Malignant neoplasm of retroperitoneum
C48.1	Malignant neoplasm of specified parts of peritoneum
C48.2	Malignant neoplasm of peritoneum, unspecified
C48.8	Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum
C49.0	Malignant neoplasm of connective and soft tissue of head, face and neck
C49.10	Malignant neoplasm of connective and soft tissue of unspecified upper limb, including shoulder
C49.11	Malignant neoplasm of connective and soft tissue of right upper limb, including shoulder

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C49.12	Malignant neoplasm of connective and soft tissue of left upper limb, including shoulder
C49.20	Malignant neoplasm of connective and soft tissue of unspecified lower limb, including hip
C49.21	Malignant neoplasm of connective and soft tissue of right lower limb, including hip
C49.22	Malignant neoplasm of connective and soft tissue of left lower limb, including hip
C49.3	Malignant neoplasm of connective and soft tissue of thorax
C49.4	Malignant neoplasm of connective and soft tissue of abdomen
C49.5	Malignant neoplasm of connective and soft tissue of pelvis
C49.6	Malignant neoplasm of connective and soft tissue of trunk, unspecified
C49.8	Malignant neoplasm of overlapping sites of connective and soft tissue
C49.9	Malignant neoplasm of connective and soft tissue, unspecified
C4A.0	Merkel cell carcinoma of lip
C4A.10	Merkel cell carcinoma of eyelid, including canthus
C4A.111	Merkel cell carcinoma of right upper eyelid, including canthus
C4A.112	Merkel cell carcinoma of right lower eyelid, including canthus
C4A.121	Merkel cell carcinoma of left upper eyelid, including canthus
C4A.122	Merkel cell carcinoma of left lower eyelid, including canthus
C4A.20	Merkel cell carcinoma of unspecified ear and external auricular canal
C4A.21	Merkel cell carcinoma of right ear and external auricular canal
C4A.22	Merkel cell carcinoma of left ear and external auricular canal
C4A.30	Merkel cell carcinoma of unspecified part of face
C4A.31	Merkel cell carcinoma of nose
C4A.39	Merkel cell carcinoma of other parts of face
C4A.4	Merkel cell carcinoma of scalp and neck
C4A.51	Merkel cell carcinoma of anal skin
C4A.52	Merkel cell carcinoma of skin of breast
C4A.59	Merkel cell carcinoma of other part of trunk
C4A.60	Merkel cell carcinoma of unspecified upper limb, including shoulder
C4A.61	Merkel cell carcinoma of right upper limb, including shoulder
C4A.62	Merkel cell carcinoma of left upper limb, including shoulder
C4A.70	Merkel cell carcinoma of unspecified lower limb, including hip
C4A.71	Merkel cell carcinoma of right lower limb, including hip
C4A.72	Merkel cell carcinoma of left lower limb, including hip
C4A.8	Merkel cell carcinoma of overlapping sites

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C4A.9	Merkel cell carcinoma, unspecified
C51.0	Malignant neoplasm of labium majus
C51.1	Malignant neoplasm of labium minus
C51.2	Malignant neoplasm of clitoris
C51.8	Malignant neoplasm of overlapping sites of vulva
C51.9	Malignant neoplasm of vulva, unspecified
C52	Malignant neoplasm of vagina
C53.0	Malignant neoplasm of endocervix
C53.1	Malignant neoplasm of exocervix
C53.8	Malignant neoplasm of overlapping sites of cervix uteri
C53.9	Malignant neoplasm of cervix uteri, unspecified
C54.0	Malignant neoplasm of isthmus uteri
C54.1	Malignant neoplasm of endometrium
C54.2	Malignant neoplasm of myometrium
C54.3	Malignant neoplasm of fundus uteri
C54.8	Malignant neoplasm of overlapping sites of corpus uteri
C54.9	Malignant neoplasm of corpus uteri, unspecified
C55	Malignant neoplasm of uterus, part unspecified
C56.1	Malignant neoplasm of right ovary
C56.2	Malignant neoplasm of left ovary
C56.3	Malignant neoplasm of bilateral ovaries
C56.9	Malignant neoplasm of unspecified ovary
C57.00	Malignant neoplasm of unspecified fallopian tube
C57.01	Malignant neoplasm of right fallopian tube
C57.02	Malignant neoplasm of left fallopian tube
C57.10	Malignant neoplasm of unspecified broad ligament
C57.11	Malignant neoplasm of right broad ligament
C57.12	Malignant neoplasm of left broad ligament
C57.20	Malignant neoplasm of unspecified round ligament
C57.21	Malignant neoplasm of right round ligament
C57.22	Malignant neoplasm of left round ligament
C57.3	Malignant neoplasm of parametrium
C57.4	Malignant neoplasm of uterine adnexa, unspecified
C57.7	Malignant neoplasm of other specified female genital organs

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C57.8	Malignant neoplasm of overlapping sites of female genital organs
C57.9	Malignant neoplasm of female genital organ, unspecified
C58	Malignant neoplasm of placenta
C61	Malignant neoplasm of prostate
C64.1	Malignant neoplasm of right kidney, except renal pelvis
C64.2	Malignant neoplasm of left kidney, except renal pelvis
C64.9	Malignant neoplasm of unspecified kidney, except renal pelvis
C65.1	Malignant neoplasm of right renal pelvis
C65.2	Malignant neoplasm of left renal pelvis
C65.9	Malignant neoplasm of unspecified renal pelvis
C66.1	Malignant neoplasm of right ureter
C66.2	Malignant neoplasm of left ureter
C66.9	Malignant neoplasm of unspecified ureter
C67.0	Malignant neoplasm of trigone of bladder
C67.1	Malignant neoplasm of dome of bladder
C67.2	Malignant neoplasm of lateral wall of bladder
C67.3	Malignant neoplasm of anterior wall of bladder
C67.4	Malignant neoplasm of posterior wall of bladder
C67.5	Malignant neoplasm of bladder neck
C67.6	Malignant neoplasm of ureteric orifice
C67.7	Malignant neoplasm of urachus
C67.8	Malignant neoplasm of overlapping sites of bladder
C67.9	Malignant neoplasm of bladder, unspecified
C68.0	Malignant neoplasm of urethra
C69.30	Malignant neoplasm of unspecified choroid
C69.31	Malignant neoplasm of right choroid
C69.32	Malignant neoplasm of left choroid
C69.40	Malignant neoplasm of unspecified ciliary body
C69.41	Malignant neoplasm of right ciliary body
C69.42	Malignant neoplasm of left ciliary body
C69.60	Malignant neoplasm of unspecified orbit
C69.61	Malignant neoplasm of right orbit

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C69.62	Malignant neoplasm of left orbit
C71.0	Malignant neoplasm of cerebrum, except lobes and ventricles
C71.1	Malignant neoplasm of frontal lobe
C71.2	Malignant neoplasm of temporal lobe
C71.3	Malignant neoplasm of parietal lobe
C71.4	Malignant neoplasm of occipital lobe
C71.5	Malignant neoplasm of cerebral ventricle
C71.6	Malignant neoplasm of cerebellum
C71.7	Malignant neoplasm of brain stem
C71.8	Malignant neoplasm of overlapping sites of brain
C71.9	Malignant neoplasm of brain, unspecified
C72.0	Malignant neoplasm of spinal cord
C72.1	Malignant neoplasm of cauda equina
C72.9	Malignant neoplasm of central nervous system, unspecified
C76.0	Malignant neoplasm of head, face and neck
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C78.6	Secondary malignant neoplasm of retroperitoneum and peritoneum
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C79.31	Secondary malignant neoplasm of brain
C7A.1	Malignant poorly differentiated neuroendocrine tumors
C7B.1	Secondary Merkel cell carcinoma
C81.10	Nodular sclerosis Hodgkin lymphoma, unspecified site
C81.11	Nodular sclerosis Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.12	Nodular sclerosis Hodgkin lymphoma, intrathoracic lymph nodes
C81.13	Nodular sclerosis Hodgkin lymphoma, intra-abdominal lymph nodes
C81.14	Nodular sclerosis Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.15	Nodular sclerosis Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.16	Nodular sclerosis Hodgkin lymphoma, intrapelvic lymph nodes
C81.17	Nodular sclerosis Hodgkin lymphoma, spleen

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

C81.18	Nodular sclerosis Hodgkin lymphoma, lymph nodes of multiple sites
C81.19	Nodular sclerosis Hodgkin lymphoma, extranodal and solid organ sites
C81.20	Mixed cellularity Hodgkin lymphoma, unspecified site
C81.21	Mixed cellularity Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.22	Mixed cellularity Hodgkin lymphoma, intrathoracic lymph nodes
C81.23	Mixed cellularity Hodgkin lymphoma, intra-abdominal lymph nodes
C81.24	Mixed cellularity Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.25	Mixed cellularity Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.26	Mixed cellularity Hodgkin lymphoma, intrapelvic lymph nodes
C81.27	Mixed cellularity Hodgkin lymphoma, spleen
C81.28	Mixed cellularity Hodgkin lymphoma, lymph nodes of multiple sites
C81.29	Mixed cellularity Hodgkin lymphoma, extranodal and solid organ sites
C81.30	Lymphocyte depleted Hodgkin lymphoma, unspecified site
C81.31	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.32	Lymphocyte depleted Hodgkin lymphoma, intrathoracic lymph nodes
C81.33	Lymphocyte depleted Hodgkin lymphoma, intra-abdominal lymph nodes
C81.34	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.35	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.36	Lymphocyte depleted Hodgkin lymphoma, intrapelvic lymph nodes
C81.37	Lymphocyte depleted Hodgkin lymphoma, spleen
C81.38	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of multiple sites
C81.39	Lymphocyte depleted Hodgkin lymphoma, extranodal and solid organ sites
C81.40	Lymphocyte-rich Hodgkin lymphoma, unspecified site
C81.41	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.42	Lymphocyte-rich Hodgkin lymphoma, intrathoracic lymph nodes
C81.43	Lymphocyte-rich Hodgkin lymphoma, intra-abdominal lymph nodes
C81.44	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.45	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.46	Lymphocyte-rich Hodgkin lymphoma, intrapelvic lymph nodes
C81.47	Lymphocyte-rich Hodgkin lymphoma, spleen
C81.48	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of multiple sites

C81.49	Lymphocyte-rich Hodgkin lymphoma, extranodal and solid organ sites
C81.70	Other Hodgkin lymphoma unspecified site
C81.71	Other Hodgkin lymphoma lymph nodes of head, face, and neck
C81.72	Other Hodgkin lymphoma intrathoracic lymph nodes
C81.73	Other Hodgkin lymphoma intra-abdominal lymph nodes
C81.74	Other Hodgkin lymphoma lymph nodes of axilla and upper limb
C81.75	Other Hodgkin lymphoma lymph nodes of inguinal region and lower limb
C81.76	Other Hodgkin lymphoma intrapelvic lymph nodes
C81.77	Other Hodgkin lymphoma spleen
C81.78	Other Hodgkin lymphoma lymph nodes of multiple sites
C81.79	Other Hodgkin lymphoma extranodal and solid organ sites
C81.90	Hodgkin lymphoma, unspecified site
C81.91	Hodgkin lymphoma, unspecified lymph nodes of head, face, and neck
C81.92	Hodgkin lymphoma, unspecified intrathoracic lymph nodes
C81.93	Hodgkin lymphoma, unspecified intra-abdominal lymph nodes
C81.94	Hodgkin lymphoma, unspecified lymph nodes of axilla and upper limb
C81.95	Hodgkin lymphoma, unspecified lymph nodes of inguinal region and lower limb
C81.96	Hodgkin lymphoma, unspecified intrapelvic lymph nodes
C81.97	Hodgkin lymphoma, unspecified spleen
C81.98	Hodgkin lymphoma, unspecified lymph nodes of multiple sites
C81.99	Hodgkin lymphoma, unspecified extranodal and solid organ sites
C83.00	Small cell B-cell lymphoma, unspecified site
C83.01	Small cell B-cell lymphoma, lymph nodes of head, face, and neck
C83.02	Small cell B-cell lymphoma, intrathoracic lymph nodes
C83.03	Small cell B-cell lymphoma, intra-abdominal lymph nodes
C83.04	Small cell B-cell lymphoma, lymph nodes of axilla and upper limb
C83.05	Small cell B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.06	Small cell B-cell lymphoma, intrapelvic lymph nodes
C83.07	Small cell B-cell lymphoma, spleen
C83.08	Small cell B-cell lymphoma, lymph nodes of multiple sites
C83.09	Small cell B-cell lymphoma, extranodal and solid organ sites

C83.30	Diffuse large B-cell lymphoma, unspecified site
C83.31	Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck
C83.32	Diffuse large B-cell lymphoma, intrathoracic lymph nodes
C83.33	Diffuse large B-cell lymphoma, intra-abdominal lymph nodes
C83.34	Diffuse large B-cell lymphoma, lymph nodes of axilla and upper limb
C83.35	Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.36	Diffuse large B-cell lymphoma, intrapelvic lymph nodes
C83.37	Diffuse large B-cell lymphoma, spleen
C83.38	Diffuse large B-cell lymphoma, lymph nodes of multiple sites
C83.39	Diffuse large B-cell lymphoma, extranodal and solid organ sites
C85.20	Mediastinal (thymic) large B-cell lymphoma, unspecified site
C85.21	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of head, face and neck
C85.22	Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes
C85.23	Mediastinal (thymic) large B-cell lymphoma, intra-abdominal lymph nodes
C85.24	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of axilla and upper limb
C85.25	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.26	Mediastinal (thymic) large B-cell lymphoma, intrapelvic lymph nodes
C85.27	Mediastinal (thymic) large B-cell lymphoma, spleen
C85.28	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of multiple sites
C85.29	Mediastinal (thymic) large B-cell lymphoma, extranodal and solid organ sites
C91.10	Chronic lymphocytic leukemia of B-cell type not having achieved remission
C91.12	Chronic lymphocytic leukemia of B-cell type in relapse
D09.0	Carcinoma in situ of bladder
D37.02	Neoplasm of uncertain behavior of tongue
D37.04	Neoplasm of uncertain behavior of the minor salivary glands
D37.05	Neoplasm of uncertain behavior of pharynx
D37.09	Neoplasm of uncertain behavior of other specified sites of the oral cavity
D37.1	Neoplasm of uncertain behavior of stomach
D37.3	Neoplasm of uncertain behavior of appendix
D37.8	Neoplasm of uncertain behavior of other specified digestive organs
D37.9	Neoplasm of uncertain behavior of digestive organ, unspecified

Medical Necessity Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

D38.0	Neoplasm of uncertain behavior of larynx
D38.5	Neoplasm of uncertain behavior of other respiratory organs
D38.6	Neoplasm of uncertain behavior of respiratory organ, unspecified
D39.2	Neoplasm of uncertain behavior of placenta
D48.60	Neoplasm of uncertain behavior of unspecified breast
D48.61	Neoplasm of uncertain behavior of right breast
D48.62	Neoplasm of uncertain behavior of left breast
O01.9	Hydatidiform mole, unspecified
Z85.00	Personal history of malignant neoplasm of unspecified digestive organ
Z85.01	Personal history of malignant neoplasm of esophagus
Z85.028	Personal history of other malignant neoplasm of stomach
Z85.038	Personal history of other malignant neoplasm of large intestine
Z85.068	Personal history of other malignant neoplasm of small intestine
Z85.09	Personal history of malignant neoplasm of other digestive organs
Z85.118	Personal history of other malignant neoplasm of bronchus and lung
Z85.12	Personal history of malignant neoplasm of trachea
Z85.22	Personal history of malignant neoplasm of nasal cavities, middle ear, and accessory sinuses
Z85.42	Personal history of malignant neoplasm of other parts of uterus
Z85.43	Personal history of malignant neoplasm of ovary
Z85.51	Personal history of malignant neoplasm of bladder
Z85.59	Personal history of malignant neoplasm of other urinary tract organ
Z85.71	Personal history of Hodgkin lymphoma
Z85.810	Personal history of malignant neoplasm of tongue
Z85.818	Personal history of malignant neoplasm of other sites of lip, oral cavity, and pharynx
Z85.820	Personal history of malignant melanoma of skin
Z85.821	Personal history of Merkel cell carcinoma
Z85.830	Personal history of malignant neoplasm of bone
Z85.831	Personal history of malignant neoplasm of soft tissue
Z85.841	Personal history of malignant neoplasm of brain
Z85.848	Personal history of malignant neoplasm of other parts of nervous tissue

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC