

# Polivy® (polatuzumab vedotin-piiq) (Intravenous)

---

**-E-**

Document Number: EOCCO-0543

Last Review Date: 06/01/2023

Date of Origin: 06/02/2020

Dates Reviewed: 06/2020, 08/2020, 01/2021, 08/2021, 08/2022, 06/2023

## I. Length of Authorization <sup>1,6</sup>

Coverage will be provided for 6 months (up to 6 cycles of therapy) and may NOT be renewed.

## II. Dosing Limits

### A. Quantity Limit (max daily dose) [NDC Unit]:

- Polivy 30 mg single-dose vial: 2 vials per 21 days
- Polivy 140 mg single-dose vial: 1 vial per 21 days

### B. Max Units (per dose and over time) [HCPCS Unit]:

- 200 billable units every 21 days

## III. Initial Approval Criteria <sup>1</sup>

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**
- Patient will receive prophylaxis for *Pneumocystis jiroveci* pneumonia and herpesvirus; **AND**
- Patient does not currently have Grade  $\geq 2$  peripheral neuropathy; **AND**
- Patient does not have CNS lymphoma; **AND**

B-Cell Lymphomas † ‡ <sup>1-5,3e</sup>

- Diffuse large B-cell lymphoma (DLBCL)  $\Phi$  or High-Grade B-Cell Lymphomas (HGBL); **AND**
  - Used as first line therapy †; **AND**
    - Used in combination with a rituximab product, cyclophosphamide, doxorubicin, and prednisone (R-CHP); **AND**
    - Patient has an International Prognostic Index (IPI) score of  $\geq 2$ ; **OR**
  - Used as subsequent treatment in patients with no intention to proceed to transplant; **AND**
    - Used in combination with rituximab OR bendamustine and rituximab; **AND**

- Used for relapsed or refractory disease >12 months after completion of first-line therapy; **OR**
- Used for primary refractory disease (partial response, no response, or progression) or relapsed disease <12 months after completion of first-line therapy in non-candidates for CAR T-cell therapy; **OR**
- Used as alternative systemic therapy (if not previously used) for relapsed/refractory disease in non-candidates for CAR T-cell therapy

**Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.**

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ◊ Orphan Drug

#### IV. Renewal Criteria <sup>1,3,4</sup>

Coverage cannot be renewed.

#### V. Dosage/Administration <sup>1,6</sup>

Indication	Dose
Previously untreated DLBCL or HGBL	Administer 1.8 mg/kg intravenously every 21 days for 6 cycles in combination with a rituximab product, cyclophosphamide, doxorubicin, and prednisone. <ul style="list-style-type: none"> <li>• Administer Polivy, cyclophosphamide, doxorubicin, and a rituximab product in any order on Day 1 after the administration of prednisone. Prednisone is administered on Days 1–5 of each cycle.</li> </ul>
Relapsed/refractory DLBCL	Administer 1.8 mg/kg intravenously every 21 days for 6 cycles in combination with bendamustine and rituximab product. <ul style="list-style-type: none"> <li>• Administer Polivy, bendamustine, and rituximab product in any order on Day 1 of each cycle.</li> </ul>
All Other Indications	Administer 1.8 mg/kg intravenously every 21 days for 6 cycles.

#### VI. Billing Code/Availability Information

HCPCS Code:

- J9309 – Injection, polatuzumab vedotin-piiq 1 mg; 1 mg = 1 billable unit

NDC(s):

- Polivy 30 mg lyophilized powder for injection, single-dose vial: 50242-0103-xx
- Polivy 140 mg lyophilized powder for injection, single-dose vial: 50242-0105-xx

## VII. References (STANDARD)

1. Polivy [package insert]. South San Francisco, CA; Genentech, Inc; April 2023. Accessed May 2023.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for polatuzumab vedotin. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2023.
3. Sehn LH, Kamdar M, Herrera AF, et al. Randomized phase 2 trial of polatuzumab vedotin (pola) with bendamustine and rituximab (BR) in relapsed/refractory (r/r) FL and DLBCL. *J Clin Oncol* 2018; 36:15\_suppl, 7507-7507.
4. Sehn LH, Herrera AF, Matasar MJ, et al. Polatuzumab vedotin (Pola) plus bendamustine (B) with rituximab (R) or obinutuzumab (G) in relapsed/refractory (R/R) Diffuse Large B-Cell Lymphoma (DLBCL): Updated results of a phase (Ph) Ib/II study (abstract). *Blood* 2018;132:Abstract 1683.
5. Tilly H, Morschhauser F, Sehn LH, et al. Polatuzumab Vedotin in Previously Untreated Diffuse Large B-Cell Lymphoma. *N Engl J Med*. 2022 Jan 27;386(4):351-363. doi: 10.1056/NEJMoa2115304.
6. Sehn LH, Herrera AF, Flowers CR, et al. Polatuzumab Vedotin in Relapsed or Refractory Diffuse Large B-Cell Lymphoma. *J Clin Oncol*. 2020 Jan 10;38(2):155-165. doi: 10.1200/JCO.19.00172.

## VIII. References (ENHANCED)

- 1e. Mounier N, El Gnaoui T, Tilly H, et al. Rituximab plus gemcitabine and oxaliplatin in patients with refractory/relapsed diffuse large B-cell lymphoma who are not candidates for high-dose therapy. A phase II Lymphoma Study Association trial. *Haematologica*. 2013;98(11):1726–1731. doi:10.3324/haematol.2013.090597.
- 2e. Morschhauser F, Flinn IW, Advani R, et al. Polatuzumab vedotin or pinatuzumab vedotin plus rituximab in patients with relapsed or refractory non-Hodgkin lymphoma: final results from a phase 2 randomised study (ROMULUS). *Lancet Haematol*. 2019 May;6(5):e254-e265. doi: 10.1016/S2352-3026(19)30026-2.
- 3e. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for B-Cell Lymphomas 3.2023. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2023.
- 4e. Magellan Health, Magellan Rx Management. Polivy Clinical Literature Review Analysis. Last updated May 2023. Accessed May 2023.

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C83.30	Diffuse large B-cell lymphoma unspecified site
C83.31	Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck
C83.32	Diffuse large B-cell lymphoma intrathoracic lymph nodes
C83.33	Diffuse large B-cell lymphoma intra-abdominal lymph nodes
C83.34	Diffuse large B-cell lymphoma lymph nodes of axilla and upper limb
C83.35	Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.36	Diffuse large B-cell lymphoma intrapelvic lymph nodes
C83.37	Diffuse large B-cell lymphoma, spleen
C83.38	Diffuse large B-cell lymphoma lymph nodes of multiple sites
C83.39	Diffuse large B-cell lymphoma extranodal and solid organ sites
C83.90	Non-follicular (diffuse) lymphoma, unspecified site
C83.91	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of head, face, and neck
C83.92	Non-follicular (diffuse) lymphoma, unspecified intrathoracic lymph nodes
C83.93	Non-follicular (diffuse) lymphoma, unspecified intra-abdominal lymph nodes
C83.94	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of axilla and upper limb
C83.95	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of inguinal region and lower limb
C83.96	Non-follicular (diffuse) lymphoma, unspecified intrapelvic lymph nodes
C83.97	Non-follicular (diffuse) lymphoma, unspecified spleen
C83.98	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of multiple sites
C83.99	Non-follicular (diffuse) lymphoma, unspecified extranodal and solid organ sites
C85.10	Unspecified B-cell lymphoma, unspecified site
C85.11	Unspecified B-cell lymphoma, lymph nodes of head, face, and neck
C85.12	Unspecified B-cell lymphoma, intrathoracic lymph nodes
C85.13	Unspecified B-cell lymphoma, intra-abdominal lymph nodes
C85.14	Unspecified B-cell lymphoma, lymph nodes of axilla and upper limb
C85.15	Unspecified B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.16	Unspecified B-cell lymphoma, intrapelvic lymph nodes
C85.17	Unspecified B-cell lymphoma, spleen
C85.18	Unspecified B-cell lymphoma, lymph nodes of multiple sites
C85.19	Unspecified B-cell lymphoma, extranodal and solid organ sites

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Articles (LCAs) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC