

Knee Cartilage Transplants

Date of Origin: 3/2005

Last Review Date: 08/28/2024

Effective Date: 9/1/2024

Dates Reviewed: 04/2013, 04/2014, 04/2015, 05/2016, 07/2016, 08/2017, 08/2019, 10/2019, 08/2020, 08/2021, 07/2022, 08/2023, 08/2024

Developed By: Medical Necessity Criteria Committee

I. Description

Allograft transplants of the knee are a type of procedure used in the treatment of individuals with symptomatic disabling cartilage injury or disease. This surgical technique can restore knee function in patients with focal articular cartilage defects due to trauma or other conditions such as osteochondritis dissecans. The procedure involves the transplantation of a piece of articular cartilage from a cadaver donor to the damaged surface of the knee.

Osteochondral autografting is a surgical procedure used to repair damaged articular cartilage. This type of procedure involves the placement of viable hyaline cartilage grafts into a cartilage defect. The grafts are harvested from a non-weight-bearing region of the joint during an open or arthroscopic procedure and then transplanted into a cartilage defect to restore the articular surface of the bone. Osteochondral autografts are performed mainly to treat small and medium-sized focal chondral and osteochondral defects of the weight-bearing surfaces of the knee joint. Two forms of osteochondral autografting are mosaicplasty and the osteochondral autograft transplantation system (OATS®) procedure. Although different instrumentation is used in mosaicplasty and OATS® procedures, the underlying principles are similar. These procedures use either multiple osteochondral cores or a single graft, harvested from a non-weight-bearing region of the joint that is autografted into the chondral defect.

Autologous Chondrocyte Transplantation (ACT) or Autologous Chondrocyte Implantation (ACI)

is a surgical procedure used to treat isolated full-thickness (down to the bone) articular cartilage defects of the knee. The first procedure is performed arthroscopically and involves harvesting a small piece of articular cartilage from the patient's knee. The cartilage biopsy is then sent to a laboratory where it is enzymatically treated to isolate chondrocytes (cartilage-producing cells of the body). The chondrocytes are then expanded in number and used later for implantation. The second stage involves arthrotomy whereby a small patch is sewn over the articular cartilage defect. The chondrocytes that have been harvested and expanded are then injected underneath this patch where they adhere to the patient's knee to form a hyaline-like cartilage that resembles the native joint cartilage.

II. Criteria: CWQI HCS-

A. Cartilage Transplants of the knee will be covered to plan limitations for **1 or more** of the following:

a. **Meniscal Allograft Transplantation** is considered medically necessary when **ALL** of the following criteria are met **CWQI: HCS-0048D**

- i. The patient has significant knee pain that interferes with age-appropriate activities of daily living and/or demands of employment
- ii. Prior significant trauma resulting in an irreparable meniscal tear or has undergone a meniscectomy where at least 50% of the meniscus has been removed
- iii. Any **ONE** of these findings following physical examination:
 1. Limited range of motion
 2. Evidence of joint swelling/effusion
 3. Joint line tenderness
- iv. Failure of provider-directed non-surgical management for at least 3 months in duration
- v. Patient has a Body Mass Index (BMI) of 35 or less
- vi. Age 49 years or younger

b. Meniscal allograft transplantation is considered NOT medically necessary for any other indication or condition, including when EITHER of the following criteria are present:

- i. Upon standing radiographs, individual demonstrates osteoarthritic changes in the knee including joint space narrowing and osteophytes which is classified by the Kellgren-Lawrence Scale as Grade III or IV
- ii. Upon MRI, an individual demonstrates articular degeneration in the affected compartment which is classified by the Modified Outerbridge Scale as Grade III or IV

c. **Osteochondral Allograft/Autograft Transplantation Systems (OATS)/Mosaicplasty**

- i. Osteochondral Allograft/Autograft Transplantation (OATS)/mosaicplasty will be covered to plan limitations when **ALL** of the following criteria have been met:
 1. The patient has significant knee pain that interferes with age-appropriate activities of daily living and/or demands of employment
 2. Presence of BOTH of the following on physical examination
 - a. A stable knee with intact or reconstructed ligaments (ACL or PCL) and menisci
 - i. Concurrent ligament stabilization or meniscal procedure at the time of OATS would be acceptable
 - b. Normal tibial-femoral and/or patella-femoral alignment
 3. Failure of provider-directed non-surgical management for at least 3 months in duration
 4. Full-thickness distal femoral articular surface (i. e. medial condyle, lateral condyle, or trochlea) and/or patellar chondral defect that has been identified during an MRI or CT arthrogram, or during an arthroscopy and classified by the Modified Outerbridge Scale as Grade III or Grade IV
 5. EITHER of the following:
 - a. Osteochondral autograft transplants and mosaicplasty:

- i. Small (less than or equal to 2.5 cm² total) chondral defects with sharp, definite borders surrounded by normal-appearing hyaline cartilage
 - b. Osteochondral allograft transplant:
 - i. Larger (less than or equal to 10.0 cm² total) chondral defects with sharp definite borders surrounded by normal appearing hyaline cartilage
- 6. Previous arthroscopic or other traditional surgical procedure (i.e., microfracture, drilling, abrasion, osteochondral graft) has resulted in an unsatisfactory outcome
- 7. Absence of inflammatory arthritis or other systemic disease affecting the joints
- 8. Minimal to absent osteoarthritic changes in the surrounding articular cartilage (e.g. Kellgren-Lawrence Grade 2 or less)
- 9. Normal articular cartilage at the lesion border (contained lesion)
- 10. For femoral and patellar chondral lesions, the absence of corresponding “kissing lesions” with a Modified Outerbridge Scale of Grade III or Grade IV of the distal femur (trochlea, condyles), patella, or tibia
- 11. The patient is not considered a candidate for total knee replacement/arthroplasty
- 12. Patient has a BMI of 35 or less
- 13. Patient is 49 years old or younger
- d. Osteochondral allograft/autograft transplantation (OATS)/mosaicplasty of the distal femoral articular or patellar surface is considered experimental and investigational for any other indication or condition.
- e. Hybrid autologous chondrocyte implantation performed with osteochondral autograft transfer system (Hybrid ACI/OATS) for the treatment of osteochondral defects is considered experimental and investigational.
- f. **Autologous Chondrocyte Transplantation (ACT) or Autologous Chondrocyte Implantation (ACI) (using the MACI™ implant) (CWQI: HCS-0048B)** will be covered to plan limitations for treatment of symptomatic single or multiple full-thickness cartilage defects of the distal femoral articular surface (i.e., medial condyle, lateral condyle or trochlea) and/or patella caused by acute or repetitive trauma when **ALL** of the following criteria are met:
 - i. The patient has significant knee pain that interferes with age-appropriate activities of daily living and/or demands of employment
 - ii. Presence of both of the following on physical examination:
 - 1. A stable knee with intact or reconstructed ligaments (ACL or PCL) and menisci
 - a. A concurrent ligament stabilization or meniscal procedure at the time of ACI would be acceptable
 - 2. Normal tibial-femoral and/or patella-femoral alignment
 - iii. Failure of provider-directed non-surgical management for at least 3 months in duration
 - iv. Full-thickness distal femoral articular surface (i.e. medial condyle, lateral condyle or trochlea) and/or patellar chondral defect of 1-10 cm² in size that has been identified during an MRI or CT arthrogram, or during arthroscopy and classified by the Modified Outerbridge Scale as Grade III or Grade IV
 - v. Absence of osteochondritis dissecans (OCD) lesion that requires bone grafting

- vi. Absence of inflammatory arthritis or other systemic disease affecting the joints
 - vii. Absence of osteoarthritis or generalized tibial chondromalacia
 - viii. Minimal to absent osteoarthritic changes in the surrounding articular cartilage (e. g. Kellgren-Lawrence Grade 2 or less)
 - ix. Normal articular cartilage at the lesion border (contained lesion)
 - x. For femoral and patellar chondral lesions, absence of a corresponding “kissing lesion” with a Modified Outerbridge Scale of Grade III or Grade IV of the distal femur (trochlea, condyles), patella or tibia
 - xi. Patient has a BMI of 35 or less
 - xii. Age 15-55 years
- g. Autologous chondrocyte implantation is considered NOT medically necessary for any other indication or condition, including when any of the following criteria are present:
- i. Any knee joint surgery within six (6) months before screening excluding surgery to procure a biopsy or a concomitant procedure to prepare the knee for a MACI implant
 - ii. Modified Outerbridge Scale Grade III or IV defect(s) on the patella or tibia
 - iii. Presence of Kellgren-Lawrence Grade 3 or 4 osteoarthritic changes in the surrounding articular cartilage
 - iv. Total meniscectomy, meniscal allograft, or bucket-handle tear or displaced tear requiring > 50% removal of the meniscus in the target knee
 - v. Septic arthritis within one (1) year before screening
 - vi. Known history of hypersensitivity to gentamicin, other aminoglycosides, or products of porcine or bovine origin
 - vii. Uncorrected congenital blood coagulation disorders
 - viii. Cruciate ligament instability
- h. Hybrid autologous chondrocyte implantation performed with an osteochondral autograft transfer system (Hybrid ACI/OATS) for the treatment of osteochondral defects is considered experimental and investigational.

***Note: The Outerbridge classification system** facilitates an objective description of chondral damage in the knee. Classifications are from a grade 0 to grade IV.

- Grade 0: Normal cartilage
- Grade I: Cartilage with swelling and softening
- Grade II: Partial thickness defect with fissures on the surface that do not reach subchondral bone or exceed 1.5cm in diameter
- Grade III: Fissuring to the level of subchondral bone in an area with a diameter greater than 1.5 cm
- Grade IV: Exposed subchondral bone.

III. Information Submitted with the Prior Authorization Request:

1. Chart notes documenting diagnosis and all current and past procedure/treatments.
2. The requested procedure description.

IV. CPT or HCPC codes covered:

Codes	Description
27407	Repair, primary, torn ligament and/or capsule, knee; cruciate
27412	Autologous chondrocyte implantation, knee
27415	Osteochondral allograft, knee, open
27416	Osteochondral autograft(s), knee, open (e.g., mosaicplasty) (includes harvesting of autograft[s]) [except to repair chondral defects of the patella] [excludes synthetic resorbable polymers]
27427	Ligamentous reconstruction (augmentation), knee; extra-articular
27428	Ligamentous reconstruction (augmentation), knee; intra-articular (open)
27429	Ligamentous reconstruction (augmentation), knee; intra-articular (open) and extra-articular
29866	Arthroscopy, knee, surgical; implantation of osteochondral autograft(s) (e.g., mosaicplasty) (includes harvesting of autografts) [except to repair chondral defects of the patella] [excludes synthetic resorbable polymers]
29867	Arthroscopy, knee, surgical; osteochondral allograft (e.g., mosaicplasty)
29868	Arthroscopy, knee, surgical; meniscal transplantation (includes arthrotomy for meniscal insertion), medial or lateral
29870	Arthroscopy, knee, diagnostic, with or without synovial biopsy (separate procedure)
J7330	Autologous cultured chondrocytes, implant
S2112	Arthroscopy, knee, surgical for harvesting of cartilage (chondrocyte cells)

V. CPT or HCPC codes NOT covered:

Codes	Description

VI. Annual Review History

Review Date	Revisions	Effective Date
04/2013	Annual Review: Added table with review date, revisions, and effective date.	04/24/2013
04/2014	Annual Review: No changes	04/30/2014
04/2015	Annual Review: No changes	04/25/2015
05/2016	Annual Review: Added criteria for OATS, and minor wording and format changes	05/26/2016
7/2016	Reformatted, Updated criteria, codes, separated into 4 separate criteria for each type to load in CWQI.	08/31/2016
08/2017	Annual Review: No changes; changed to new template	08/23/2017
08/2019	Annual Review: Reformatted and updated criteria, to provide clarity on requirements for coverage and removed duplicated language	09/01/2019
10/2019	Update: Corrected c.i.5.b.i. 'greater than or equal to 10.0 cm ² total' to ensure clarity of the requirement	10/09/2019
08/2020	Annual Review: No changes	09/01/2020
08/2021	Annual Review: No changes	09/01/2021
07/2022	Annual Review: updates to align with eviCore, added menisci, Absence of osteoarthritis or generalized tibial chondromalacia	08/01/2022
08/2023	Annual Review: updates to align with evCore- added 'A concurrent ligament stabilization or meniscal procedure at the time of ACI would be acceptable'	09/01/2023
08/2024	Annual Review: No changes	09/01/2024
01/2025	Updated the imaging findings for osteochondral allograft transplants also required to be less than or equal to 10.0cm ² total	

VII. References

1. American Academy of Orthopedic Surgeons. Advisory Statement: Use of musculoskeletal tissue allografts. February 2001; Document No. 1011.
2. Bugbee WD. Fresh osteochondral allografts. Journal of Knee Surgery. Summer 2002; 15(3):191-5.

3. Caldwell PE 3rd, Sherton WR. Indications for allografts. *Orthop Clin North Am.* 2005 Oct; 36(4):459-67.
4. Detterline AJ, Goldberg S, Bach BR Jr, Cole BJ. Treatment options for articular cartilage defects of the knee. *Orthop Nurs.* 2005 Sep-Oct; 24(5):361-6.
5. Dopirak RM, Steensen RN, Maurus PB. The medial patellofemoral ligament. *Orthopedics.* 2008;31(4):331-338.
6. Felix NA, Paulos LE. Current status of meniscal transplantation. *Knee.* 2003;10(1):13-17.
7. Giannini S, Buda R, Grigolo B, et al. Bipolar fresh osteochondral allograft of the ankle. *Foot Ankle Int.* 2010;31(1):38-46.
8. Gross AE, Kim W, Las Heras F, et al. Fresh osteochondral allografts for posttraumatic knee defects: Long-term follow up. *Clin Orthop Relat Res.* 2008;466(8):1863-1870.
9. Gross AE. Repair of cartilage defects in the knee. *Journal of Knee Surgery.* Summer 2002; 15(3):167-9.
10. Hangody L, Fules P. Autologous osteochondral mosaicplasty for the treatment of full-thickness defects of weight-bearing joints. *The Journal of bone and Joint Surgery.* 2003; 85:25-32
11. Krych AJ, Jackson JD, Hoskin TL, Dahm DL. A meta-analysis of patellar tendon autograft versus patellar tendon allograft in anterior cruciate ligament reconstruction. *Arthroscopy* 2008;24(3):292-298.
12. Ma HL, Hung SC, Wang ST, et al. Osteochondral autografts transfer for post-traumatic osteochondral defect of the knee-2 to 5 years follow-up. *Injury.* 2004 Dec; 35(12):1286-92.
13. Melton JT, Wilson AJ, Chapman-Sheath P, Cossey AJ. TruFit CB bone plug: Chondral repair, scaffold design, surgical technique, and early experiences. *Expert Rev Med Devices.* 2010;7(3):333-341
14. Noyes FR, Barber-Westin SD, Rankin M. Meniscal transplantation in symptomatic patients less than fifty years old. *J. Bone Joint Surg Am.* 2004 July; 86-A (7):1392-404.
15. Prodromos C, Joyce B, Shi K. A meta-analysis of stability of autografts compared to allografts after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(7):851-856.
16. Rose T, Craatz S, Hepp P, et al. The autologous osteochondral transplantation of the knee: clinical results, radiographic findings and histological aspects. *Arch Orthop Trauma Surg.* 2005 Nov; 125(9):628-37.
17. Sekiya JK, West RV, Groff YJ, et al. Clinical outcomes following isolated lateral meniscal allograft transplantation. *Arthroscopy.* 2006 Jul; 22(7):771-80.
18. Shasha N, Aubin PP, Cheah HK, et al. Long-term clinical experience with fresh osteochondral allografts for articular knee defects in high demand patients. *Cell Tissue Bank.* 2002; 3(3):175-82.
19. Tins BJ, McCall IW, Takahashi T, et al. Autologous chondrocyte implantation in knee joint: MR imaging and histologic features at 1-year follow-up. *Radiology.* 2005 Feb; 234(2):501-8.
20. Van Arkel E, de Boer HH. Human meniscal transplantation: Preliminary results at 2 to 5 year follow-up. *J Bone Joint Surg.* 1995;77(4):589-595.
21. Wilcox T, Goble EM. Indications for meniscal allograft reconstruction. *Am J Knee Surg.* 1996; 9:35-36.
22. Washington University Physicians. Autologous chondrocyte implantation
<https://www.ortho.wustl.edu/content/Patient-Care/2888/Services/Sports-Medicine/Overview/Knee/Autologous-Chondrocyte-Implantation.aspx>
23. Physician Advisors

Appendix 1 – Applicable Diagnosis Codes:

Codes	Description
M22.2X1 - M22.3X9	Patellofemoral disorders and other derangements of patella [including lateral, medial, anterior and posterior ligaments]
M22.8X1 - M22.8X9	Other disorders of patella [including lateral, medial, anterior and posterior ligaments]
M23.00-M23.92	Internal derangement of knee [articular cartilage defect]
M23.601 - M23.8X9	Other spontaneous disruption of ligament(s) of knee and other internal derangements of knee [including lateral, medial, anterior and posterior ligaments]
M25.861-M25.869	Other specified joint disorders, knee [articular cartilage of knee]
M93.261-M93.269	Osteochondritis of knee
Codes	Description
M22.2X1 - M22.3X9	Patellofemoral disorders and other derangements of patella [including lateral, medial, anterior and posterior ligaments]
M22.8X1 - M22.8X9	Other disorders of patella [including lateral, medial, anterior and posterior ligaments]
M23.00-M23.92	Internal derangement of knee [articular cartilage defect]

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD):

Jurisdiction(s): 5, 8	NCD/LCD Document (s):
Noridian Local Coverage Determination (LCD) Non-covered Services (L35008) – CPT 28446	
https://med.noridianmedicare.com/documents/10546/6990983/Non-Covered+Services+LCD	

NCD/LCD Document (s):**Medicare Part B Administrative Contractor (MAC) Jurisdictions**

Jurisdiction	Applicable State/US Territory	Contractor
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC